הודעה על החמרה (מידע בטיחות) בעלון לרופא (מעודכן 05.2013)

תאריך <u>25/07/2017</u>

שם תכשיר באנגלית ומספר הרישום [131-59-31060-00] שם בעל הרישום: Roche Pharmaceuticals (Israel) Ltd

! טופס זה מיועד לפרוט ההחמרות בלבד

טופט זה מיועד לפרוט ההוומדות בלבר ! ההחמרות המבוקשות - עלון לרופא			
פרק בעלון טקסט נוכחי טקסט חדש			
[]	[]	4.8 Undesirable effects	
Table 1: Adverse Reactions by Frequency	Table 1: Adverse Reactions by Frequency		
Metabolism and nutrition disorders: Hypomagnesaemia and Hyponatraemia were added as a "Very Common" adverse reactions.	בעדכון עלון זה ישנה הוספת מידע חדש		
Respiratory, thoracic and mediastinal disorders: Epistaxis and Cough were added as a "Very Common" adverse reactions.	common מופיע בעמודת Epistaxis		
Musculoskeletal and connective	common מופיע בעמודת Myalgia		
tissue disorders: Myalgia was added as a "Very Common" adverse reaction. []	[]		
Table 2: Severe Adverse Reactions by Frequency	Table 2: Severe Adverse Reactions by Frequency		
Metabolism and nutrition disorders: Hyponatraemia was added as a "Common" adverse reaction.	בעדכון עלון זה ישנה הוספת מידע חדש		
[]	[]		
<u>Description of selected serious</u> <u>adverse reactions</u>	<u>Description of selected serious</u> <u>adverse reactions</u>		
GI-vaginal Fistulae in study GOG- 0240 []	GI-vaginal Fistulae in study GOG- 0240 []		
The frequency of GI-vaginal fistulae in the group treated with Avastin + chemotherapy was higher in patients with recurrence within the field of prior radiation (16.7%) compared with patients with no prior radiation and/ or no	The frequency of GI-vaginal fistulae in the group treated with Avastin + chemotherapy was higher in patients with recurrence within the field of prior radiation (16.7%) compared with patients with recurrence outside the field		

ההחמרות המבוקשות - עלון לרופא			
טקסט חדש	טקסט נוכחי	פרק בעלון	
recurrence inside the field of prior radiation (3.6%). []	of prior radiation (3.6%). []		
Wound healing [] In clinical trials of ovarian cancer, Grade 3-5 wound healing complications were observed in up to 1.8% of patients in the bevacizumab arm versus 0.1% in the control arm (NCI-CTCAE v.3). []	Wound healing [] In clinical trials of ovarian cancer, Grade 3-5 wound healing complications were observed in up to 1.2% of patients in the bevacizumab arm versus 0.1% in the control arm (NCI-CTCAE v.3). []		
Proteinuria [] Proteinuria ranged in severity from clinically asymptomatic, transient, trace proteinuria to nephrotic syndrome, with the great majority as Grade 1 proteinuria (NCI-CTCAE v.3). Grade 3 proteinuria was reported in up to 10.9% of treated patients. []	Proteinuria [] Proteinuria ranged in severity from clinically asymptomatic, transient, trace proteinuria to nephrotic syndrome, with the great majority as Grade 1 proteinuria (NCI-CTCAE v.3). Grade 3 proteinuria was reported in up to 8.1% of treated patients. []		

מצ"ב העלון, שבו מסומנות ההחמרות המבוקשות <mark>על רקע צהוב</mark>.

שינויים שאינם בגדר החמרות סומנו <u>(בעלוו)</u> בצבע שונה. יש לסמן רק תוכן מהותי ולא שינויים במיקום הטקסט.