

אוגוסט 2022

רופא/ה נכבד/ה,

רוקח/ת נכבד/ה,

## RINVOQ

Prolonged release tablets

(Upadacitinib 15mg, 30mg as hemihydrate)

חברת **AbbVie Biopharmaceuticals Ltd.** שמחה להודיע על אישור התוויה חדשה לתכשיר רינבוק לטיפול בקוליטיס כיבית.

נוסח ההתוויה החדש כפי שאושרה על ידי משרד הבריאות:

### Ulcerative Colitis

**RINVOQ is indicated for the treatment of adult patients with moderately to severely active ulcerative colitis who have had an inadequate response, lost response or were intolerant to either conventional therapy or a biologic agent.**

לרינבוק תוכנית ניהול סיכונים מותאמת לישראל המאושרת על ידי משרד הבריאות.

להודעה זו מצורפים עלונים עם סימון השינויים, כרטיס מידע בטיחותי למטופל וחוברת לצוות הרפואי. בעלונים, מידע שהתווסף מסומן באדום ומידע שהוסר מסומן בכחול.

העלונים לרופא ולצרכן נשלחו למאגר התרופות שבאתר משרד הבריאות, וניתן לקבלם מודפסים על ידי פניה לבעל הרישום, AbbVie Biopharmaceuticals Ltd, רחוב החרש 4, הוד השרון או בטלפון 7909600 – 09.

בברכה,

רחלי אברמוביץ'  
רוקחת ממונה

## **RINVOQ®**

### **1. NAME OF THE MEDICINAL PRODUCT**

RINVOQ 15 mg prolonged-release tablets  
RINVOQ 30 mg prolonged-release tablets

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

#### **RINVOQ 15 mg prolonged-release tablets**

Each prolonged-release tablet contains upadacitinib hemihydrate, equivalent to 15 mg of upadacitinib.

#### **RINVOQ 30 mg prolonged-release tablets**

Each prolonged-release tablet contains upadacitinib hemihydrate, equivalent to 30 mg of upadacitinib.

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Prolonged-release tablet

#### **RINVOQ 15 mg prolonged-release tablets**

Purple 14 x 8 mm, oblong biconvex prolonged-release tablets imprinted on one side with 'a15'.

#### **RINVOQ 30 mg prolonged-release tablets**

Red 14 x 8 mm, oblong biconvex prolonged-release tablets imprinted on one side with 'a30'.

### **4. CLINICAL PARTICULARS**

#### **Patient safety information Card**

The marketing of RINVOQ is subject to a risk management plan (RMP) including a 'Patient safety information card'. The 'Patient safety information card', emphasizes important safety information that the patient should be aware of before and during treatment. Please explain to the patient the need to review the card before starting treatment.

#### **HCP educational brochure**

This product is marketed with HCP educational brochure providing important safety information. Please ensure you are familiar with this material as it contains important safety information.

#### **4.1 Therapeutic indications**

##### **Rheumatoid arthritis**

RINVOQ is indicated for the treatment of moderate to severe active rheumatoid arthritis in adult patients who have responded inadequately to, or who are intolerant to one or more disease-modifying anti-rheumatic drugs (DMARDs). RINVOQ may be used as monotherapy or in combination with methotrexate.

### Psoriatic arthritis

RINVOQ is indicated for the treatment of active psoriatic arthritis in adult patients who have responded inadequately to, or who are intolerant to one or more DMARDs. RINVOQ may be used as monotherapy or in combination with methotrexate.

### Ankylosing spondylitis

RINVOQ is indicated for the treatment of active ankylosing spondylitis in adult patients who have responded inadequately to conventional therapy.

### Atopic dermatitis

RINVOQ is indicated for the treatment of moderate to severe atopic dermatitis in adults and adolescents 12 years and older who are candidates for systemic therapy.

### Ulcerative colitis

RINVOQ is indicated for the treatment of adult patients with moderately to severely active ulcerative colitis who have had an inadequate response, lost response or were intolerant to either conventional therapy or a biologic agent.

## **4.2 Posology and method of administration**

Treatment with upadacitinib should be initiated and supervised by physicians experienced in the diagnosis and treatment of conditions for which upadacitinib is indicated.

### Posology

#### Rheumatoid arthritis, psoriatic arthritis and ankylosing spondylitis

The recommended dose of upadacitinib is 15 mg once daily.

Consideration should be given to discontinuing treatment in patients with ankylosing spondylitis who have shown no clinical response after 16 weeks of treatment. Some patients with initial partial response may subsequently improve with continued treatment beyond 16 weeks.

### Atopic dermatitis

#### Adults

The recommended dose of upadacitinib is 15 mg or 30 mg once daily based on individual patient presentation.

- A dose of 30 mg once daily may be appropriate for patients with high disease burden.
- A dose of 30 mg once daily may be appropriate for patients with an inadequate response to 15 mg once daily.
- The lowest effective dose for maintenance should be considered.

For patients  $\geq$  65 years of age, the recommended dose is 15 mg once daily.

#### Adolescents (from 12 to 17 years of age)

The recommended dose of upadacitinib is 15 mg once daily for adolescents weighing at least 30 kg.

#### Concomitant topical therapies

Upadacitinib can be used with or without topical corticosteroids. Topical calcineurin inhibitors may be used for sensitive areas such as the face, neck, and intertriginous and genital areas.

Consideration should be given to discontinuing upadacitinib treatment in any patient who shows no evidence of therapeutic benefit after 12 weeks of treatment.

### Ulcerative colitis

#### Induction

The recommended induction dose of upadacitinib is 45 mg once daily (three tablets of 15 mg) for 8 weeks. For patients who do not achieve adequate therapeutic benefit by week 8, upadacitinib 45 mg once daily may be continued for an additional 8 weeks (see sections 4.8 and 5.1). Upadacitinib should be discontinued in any patient who shows no evidence of therapeutic benefit by week 16.

#### Maintenance

The recommended maintenance dose of upadacitinib is 15 mg (one tablet of 15 mg) or 30 mg (one tablet of 30 mg or two tablets of 15 mg) once daily based on individual patient presentation:

- A dose of 30 mg once daily may be appropriate for some patients, such as those with high disease burden or requiring 16-week induction treatment.
- A dose of 30 mg once daily may be appropriate for patients who do not show adequate therapeutic benefit to 15 mg once daily.
- The lowest effective dose for maintenance should be considered.

For patients  $\geq 65$  years of age, the recommended dose is 15 mg once daily.

In patients who have responded to treatment with upadacitinib, corticosteroids may be reduced and/or discontinued in accordance with standard of care.

#### Interactions

For patients with ulcerative colitis receiving strong inhibitors of cytochrome P450 (CYP) 3A4 (e.g., ketoconazole, clarithromycin), the recommended induction dose is 30 mg once daily and the recommended maintenance dose is 15 mg once daily (see section 4.5).

#### Dose initiation

Treatment should not be initiated in patients with an absolute lymphocyte count (ALC) that is  $< 0.5 \times 10^9$  cells/L, an absolute neutrophil count (ANC) that is  $< 1 \times 10^9$  cells/L or who have haemoglobin (Hb) levels that are  $< 8$  g/dL (see sections 4.4 and 4.8).

#### *Dose interruption*

Treatment should be interrupted if a patient develops a serious infection until the infection is controlled.

Interruption of dosing may be needed for management of laboratory abnormalities as described in Table 1.

#### **Table 1. Laboratory measures and monitoring guidance**

Laboratory measure	Action	Monitoring guidance
Absolute Neutrophil Count (ANC)	Treatment should be interrupted if ANC is $< 1 \times 10^9$ cells/L and may be restarted once ANC returns above this value	Evaluate at baseline and then no later than 12 weeks after initiation of treatment. Thereafter evaluate according to individual patient management.
Absolute Lymphocyte Count (ALC)	Treatment should be interrupted if ALC is $< 0.5 \times 10^9$ cells/L and may be restarted once ALC returns above this value	
Haemoglobin (Hb)	Treatment should be interrupted if Hb is $< 8$ g/dL and may be restarted once Hb returns above this value	
Hepatic transaminases	Treatment should be temporarily interrupted if drug-induced liver injury is suspected	Evaluate at baseline and thereafter according to routine patient management.
Lipids	Patients should be managed according to international clinical guidelines for hyperlipidaemia	<b>Evaluate</b> 12 weeks after initiation of treatment and thereafter according to international clinical guidelines for hyperlipidaemia

### Special populations

#### *Elderly*

#### *Rheumatoid arthritis, psoriatic arthritis, ankylosing spondylitis*

There are limited data in patients aged 75 years and older.

#### *Atopic dermatitis*

For atopic dermatitis, doses higher than 15 mg once daily are not recommended in patients aged 65 years and older (see [Section-section 4.8](#)).

~~There are limited data in patients aged 75 years and older.~~

#### *Ulcerative colitis*

For ulcerative colitis, doses higher than 15 mg once daily for maintenance therapy are not recommended in patients aged 65 years and older (see section 4.8). The safety and efficacy of upadacitinib in patients aged 75 and older have not yet been established.

#### *Renal impairment*

No dose adjustment is required in patients with mild or moderate renal impairment. There are limited data on the use of upadacitinib in subjects with severe renal impairment (see section 5.2). Upadacitinib ~~15 mg once daily~~ was not evaluated in clinical trials in patients with eGFR<40 mL/min/1.73 m<sup>2</sup>. Upadacitinib should be used with caution in patients with severe renal impairment as described in Table 2. ~~Upadacitinib 30 mg once daily is not recommended for patients with severe renal impairment.~~ The use of upadacitinib has not been studied in subjects with end stage renal disease and is therefore not recommended for use in these patients.

**Table 2 Recommended dose for severe renal impairment<sup>a</sup>**

<b>Therapeutic indication</b>	<b>Recommended once daily dose</b>
<u>Rheumatoid arthritis, psoriatic arthritis, ankylosing spondylitis, atopic dermatitis</u>	<u>15 mg</u>
<u>Ulcerative colitis</u>	<u>Induction: 30 mg</u>
	<u>Maintenance: 15 mg</u>
<u><sup>a</sup>estimated glomerular filtration rate (eGFR) 15 to &lt; 30 ml/min/1.73m<sup>2</sup></u>	

#### *Hepatic impairment*

No dose adjustment is required in patients with mild (Child -Pugh A) or moderate (Child -Pugh B) hepatic impairment (see section 5.2). Upadacitinib should not be used in patients with severe (Child -Pugh C) hepatic impairment (see section 4.3).

#### *Paediatric population*

The safety and efficacy of RINVOQ in children with atopic dermatitis below the age of 12 years have not been established. No data are available. No clinical exposure data are available in adolescents < 40 kg (see section 5.2).

The safety and efficacy of RINVOQ in children and adolescents with rheumatoid arthritis, psoriatic arthritis, ~~and~~ ankylosing spondylitis and ulcerative colitis aged 0 to less than 18 years have not yet been established. No data are available.

#### Method of administration

RINVOQ is to be taken orally once daily with or without food and may be taken at any time of the day. Tablets should be swallowed whole and should not be split, crushed, or chewed in order to ensure the entire dose is delivered correctly.

#### **4.3 Contraindications**

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Active tuberculosis (TB) or active serious infections (see section 4.4).
- Severe hepatic impairment (see section 4.2).
- Pregnancy (see section 4.6).

#### **4.4 Special warnings and precautions for use**

##### Immunosuppressive medicinal products

Combination with other potent immunosuppressants such as azathioprine, 6-mercaptopurine, ciclosporin, tacrolimus, and biologic DMARDs or other Janus kinase (JAK) inhibitors has not been

evaluated in clinical studies and is not recommended as a risk of additive immunosuppression cannot be excluded.

### Serious infections

Serious and sometimes fatal infections have been reported in patients receiving upadacitinib. The most frequent serious infections reported with upadacitinib included pneumonia and cellulitis (see section 4.8). Cases of bacterial meningitis have been reported in patients receiving upadacitinib. Among opportunistic infections, tuberculosis, multidermatomal herpes zoster, oral/oesophageal candidiasis, and cryptococcosis were reported with upadacitinib.

Upadacitinib should not be initiated in patients with an active, serious infection, including localised infections.

Consider the risks and benefits of treatment prior to initiating upadacitinib in patients:

- with chronic or recurrent infection
- who have been exposed to tuberculosis
- with a history of a serious or an opportunistic infection
- who have resided or travelled in areas of endemic tuberculosis or endemic mycoses; or
- with underlying conditions that may predispose them to infection.

Patients should be closely monitored for the development of signs and symptoms of infection during and after treatment with upadacitinib. Upadacitinib therapy should be interrupted if a patient develops a serious or opportunistic infection. A patient who develops a new infection during treatment with upadacitinib should undergo prompt and complete diagnostic testing appropriate for an immunocompromised patient; appropriate antimicrobial therapy should be initiated, the patient should be closely monitored, and upadacitinib therapy should be interrupted if the patient is not responding to antimicrobial therapy. Upadacitinib therapy may be resumed once the infection is controlled.

As there is a higher incidence of infections in the elderly  $\geq 65$  years of age, caution should be used when treating this population.

### *Tuberculosis*

Patients should be screened for tuberculosis (TB) before starting upadacitinib therapy. Upadacitinib should not be given to patients with active TB (see section 4.3). Anti-TB therapy should be considered prior to initiation of upadacitinib in patients with previously untreated latent TB or in patients with risk factors for TB infection.

Consultation with a physician with expertise in the treatment of TB is recommended to aid in the decision about whether initiating anti-TB therapy is appropriate for an individual patient.

Patients should be monitored for the development of signs and symptoms of TB, including patients who tested negative for latent TB infection prior to initiating therapy.

### Viral reactivation

Viral reactivation, including cases of herpes virus reactivation (e.g., herpes zoster), was reported in clinical studies (see section 4.8). The risk of herpes zoster appears to be higher in Japanese patients treated with upadacitinib. If a patient develops herpes zoster, interruption of upadacitinib therapy should be considered until the episode resolves.

Screening for viral hepatitis and monitoring for reactivation should be performed before starting and during therapy with upadacitinib. Patients who were positive for hepatitis C antibody and hepatitis C virus RNA were excluded from clinical studies. Patients who were positive for hepatitis B surface

antigen or hepatitis B virus DNA were excluded from clinical studies. If hepatitis B virus DNA is detected while receiving upadacitinib, a liver specialist should be consulted.

### Vaccination

No data are available on the response to vaccination with live vaccines in patients receiving upadacitinib. Use of live, attenuated vaccines during or immediately prior to upadacitinib therapy is not recommended. Prior to initiating upadacitinib, it is recommended that patients be brought up to date with all immunisations, including prophylactic zoster vaccinations, in agreement with current immunisation guidelines. (see section 5.1 for data on inactivated pneumococcal polysaccharide conjugate vaccine (13-valent, adsorbed) and concomitant use with upadacitinib).

### Malignancy

The risk of malignancies, including lymphoma is increased in patients with rheumatoid arthritis. Immunomodulatory medicinal products may increase the risk of malignancies, including lymphoma. The clinical data are currently limited and long-term studies are ongoing.

Malignancies were observed in clinical studies of upadacitinib. The risks and benefits of upadacitinib treatment should be considered prior to initiating therapy in patients with a known malignancy other than a successfully treated non-melanoma skin cancer (NMSC) or when considering continuing upadacitinib therapy in patients who develop a malignancy.

#### *Non-melanoma skin cancer*

NMSCs have been reported in patients treated with upadacitinib. Periodic skin examination is recommended for patients who are at increased risk for skin cancer.

### Haematological abnormalities

Absolute Neutrophil Count (ANC)  $< 1 \times 10^9$  cells/L, Absolute Lymphocyte Count (ALC)  $< 0.5 \times 10^9$  cells/L and haemoglobin  $< 8$  g/dL were reported in  $\leq 1$  % of patients in clinical trials (see section 4.8). Treatment should not be initiated, or should be temporarily interrupted, in patients with an ANC  $< 1 \times 10^9$  cells/L, ALC  $< 0.5 \times 10^9$  cells/L or haemoglobin  $< 8$  g/dL observed during routine patient management (see section 4.2).

### Diverticulitis

Events of diverticulitis have been reported in clinical trials and from post-marketing sources. Diverticulitis may cause gastrointestinal perforation. Upadacitinib should be used with caution in patients with diverticular disease and especially in patients chronically treated with concomitant medications associated with an increased risk of diverticulitis: nonsteroidal anti-inflammatory drugs, corticosteroids, and opioids. Patients presenting with new onset abdominal signs and symptoms should be evaluated promptly for early identification of diverticulitis to prevent gastrointestinal perforation

### Cardiovascular risk

Rheumatoid arthritis patients have an increased risk for cardiovascular disorders. Patients treated with upadacitinib should have risk factors (e.g., hypertension, hyperlipidaemia) managed as part of usual standard of care.

### Lipids

Treatment with upadacitinib was associated with dose-dependent increases in lipid parameters, including total cholesterol, low-density lipoprotein (LDL) cholesterol, and high-density lipoprotein (HDL) cholesterol (see section 4.8). Elevations in LDL cholesterol decreased to pre-treatment levels in response to statin therapy, although evidence is limited. The effect of these lipid parameter elevations



on cardiovascular morbidity and mortality has not been determined (see section 4.2 for monitoring guidance).

#### Hepatic transaminase elevations

Treatment with upadacitinib was associated with an increased incidence of liver enzyme elevation compared to placebo.

Evaluate at baseline and thereafter according to routine patient management. Prompt investigation of the cause of liver enzyme elevation is recommended to identify potential cases of drug-induced liver injury.

If increases in ALT or AST are observed during routine patient management and drug-induced liver injury is suspected, upadacitinib therapy should be interrupted until this diagnosis is excluded.

#### Venous thromboembolism

Events of deep venous thrombosis (DVT) and pulmonary embolism (PE) have been reported in patients receiving JAK inhibitors including upadacitinib. Upadacitinib should be used with caution in patients at high risk for DVT/PE. Risk factors that should be considered in determining the patient's risk for DVT/PE include older age, obesity, a medical history of DVT/PE, patients undergoing major surgery, and prolonged immobilisation. If clinical features of DVT/PE occur, upadacitinib treatment should be discontinued and patients should be evaluated promptly, followed by appropriate treatment.

#### Elderly

There is an increased risk of adverse reactions with the upadacitinib dose of 30 mg once daily in patients aged 65 years and older. The recommended dose for long-term use is 15 mg once daily for this patient population (see sections 4.2 and 4.8).

### **4.5 Interaction with other medicinal products and other forms of interaction**

#### Potential for other medicinal products to affect the pharmacokinetics of upadacitinib

Upadacitinib is metabolised mainly by CYP3A4. Therefore, upadacitinib plasma exposures can be affected by medicinal products that strongly inhibit or induce CYP3A4.

#### *Coadministration with CYP3A4 inhibitors*

Upadacitinib exposure is increased when co-administered with strong CYP3A4 inhibitors (such as ketoconazole, itraconazole, posaconazole, voriconazole, clarithromycin, and grapefruit). In a clinical study, coadministration of upadacitinib with ketoconazole resulted in 70% and 75% increases in upadacitinib  $C_{max}$  and AUC, respectively. Upadacitinib 15 mg once daily should be used with caution in patients receiving chronic treatment with strong CYP3A4 inhibitors. Upadacitinib 30 mg once daily dose is not recommended for patients with atopic dermatitis receiving chronic treatment with strong CYP3A4 inhibitors. For patients with ulcerative colitis using strong CYP3A4 inhibitors, the recommended induction dose is 30 mg once daily (for up to 16 weeks) and the recommended maintenance dose is 15 mg once daily (see section 4.2). Alternatives to strong CYP3A4 inhibitor medications should be considered when used in the long-term. Food or drink containing grapefruit should be avoided during treatment with upadacitinib.

#### *Coadministration with CYP3A4 inducers*

Upadacitinib exposure is decreased when co-administered with strong CYP3A4 inducers (such as rifampin and phenytoin), which may lead to reduced therapeutic effect of upadacitinib. In a clinical study, co-administration of upadacitinib after multiple doses of rifampicin (strong CYP3A inducer)

resulted in approximately 50% and 60% decreases in upadacitinib  $C_{max}$  and AUC, respectively. Patients should be monitored for changes in disease activity if upadacitinib is co-administered with strong CYP3A4 inducers.

Methotrexate and pH modifying medicinal products (e.g., antacids or proton pump inhibitors) have no effect on upadacitinib plasma exposures.

#### Potential for upadacitinib to affect the pharmacokinetics of other medicinal products

Administration of multiple 30 mg or 45 mg once daily doses of upadacitinib to healthy subjects had a limited effect on midazolam (sensitive substrate for CYP3A) plasma exposures (24-26% decrease in midazolam AUC and  $C_{max}$ ), indicating that upadacitinib 30 mg or 45 mg once daily may have a weak induction effect on CYP3A. In a clinical study, rosuvastatin and atorvastatin AUC were decreased by 33% and 23%, respectively, and rosuvastatin  $C_{max}$  was decreased by 23% following the administration of multiple 30 mg once daily doses of upadacitinib to healthy subjects. Upadacitinib had no relevant effect on atorvastatin  $C_{max}$  or on plasma exposures of ortho-hydroxyatorvastatin (major active metabolite for atorvastatin). Administration of multiple 45 mg once daily doses of upadacitinib to healthy subjects led to a limited increase in AUC and  $C_{max}$  of dextromethorphan (sensitive CYP2D6 substrate) by 30% and 35%, respectively, indicating that upadacitinib 45 mg once daily has a weak inhibitory effect on CYP2D6. No dose adjustment is recommended for CYP3A substrates, CYP2D6 substrates, ~~or for~~ rosuvastatin or atorvastatin when coadministered with upadacitinib.

Upadacitinib has no relevant effects on plasma exposures of ethinylestradiol, levonorgestrel, methotrexate, or medicinal products that are substrates for metabolism by CYP1A2, CYP2B6, CYP2C9, or CYP2C19, ~~or~~ CYP2D6.

## 4.6 Fertility, pregnancy and lactation

### Women of childbearing potential

Women of childbearing potential should be advised to use effective contraception during treatment and for 4 weeks following the final dose of upadacitinib. Female paediatric patients and/or their parents/caregivers should be informed about the need to contact the treating physician once the patient experiences menarche while taking upadacitinib.

### Pregnancy

There are no or limited data on the use of upadacitinib in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3). Upadacitinib was teratogenic in rats and rabbits with effects in bones in rat foetuses and in the heart in rabbit foetuses when exposed *in utero*.

Upadacitinib is contraindicated during pregnancy (see section 4.3).

If a patient becomes pregnant while taking upadacitinib the parents should be informed of the potential risk to the foetus.

### Breast-feeding

It is unknown whether upadacitinib/metabolites are excreted in human milk. Available pharmacodynamic/toxicological data in animals have shown excretion of upadacitinib in milk (see section 5.3).

A risk to newborns/infants cannot be excluded.

Upadacitinib should not be used during breast-feeding. A decision must be made whether to discontinue breast-feeding or to discontinue upadacitinib therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

## Fertility

The effect of upadacitinib on human fertility has not been evaluated. Animal studies do not indicate effects with respect to fertility (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

Upadacitinib has no or negligible influence on the ability to drive and use machines.

### **4.8 Undesirable effects**

#### Summary of the safety profile

In the placebo-controlled clinical trials for rheumatoid arthritis, psoriatic arthritis, and ankylosing spondylitis, the most commonly reported adverse reactions ( $\geq 2\%$  of patients in at least one of the indications with the highest rate among indications presented) with upadacitinib 15 mg were upper respiratory tract infections (19.5%), blood creatine phosphokinase (CPK) increased (8.6%), alanine transaminase increased (4.3%), bronchitis (3.9%), nausea (3.5%), neutropaenia (2.8%), cough (2.2%), aspartate transaminase increased (2.2%), and hypercholesterolaemia (2.2%).

In the placebo-controlled atopic dermatitis clinical trials, the most commonly reported adverse reactions ( $\geq 2\%$  of patients) with upadacitinib 15 mg or 30 mg were upper respiratory tract infection (25.4%), acne (15.1%), herpes simplex (8.4%), headache (6.3%), CPK increased (5.5%), cough (3.2%), folliculitis (3.2%), abdominal pain (2.9%), nausea (2.7%), neutropaenia (2.3%), pyrexia (2.1%), and influenza (2.1%).

The most common serious adverse reactions were serious infections (see section 4.4).

The safety profile of upadacitinib with long-term treatment was generally similar to the safety profile during the placebo-controlled period across indications.

In the placebo-controlled ulcerative colitis induction and maintenance clinical trials, the most commonly reported adverse reactions ( $\geq 3\%$  of patients) with upadacitinib 45 mg, 30 mg or 15 mg were upper respiratory tract infection (19.9%), blood CPK increased (7.6%), acne (6.3%), neutropaenia (6.0%), rash (5.2%), herpes zoster (4.4%), hypercholesterolemia (4.0%), folliculitis (3.6%), herpes simplex (3.2%), and influenza (3.2%).

#### Tabulated list of adverse reactions

The following list of adverse reactions is based on experience from clinical studies.

The frequency of adverse reactions listed below is defined using the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ). The frequencies in Table 32 are based on the higher of the rates for adverse reactions reported with RINVOQ 15 mg in clinical trials of rheumatologic disease (15 mg), and atopic dermatitis (15 mg and 30 mg) or ulcerative colitis (15 mg, 30 mg and 45 mg) clinical trials, or RINVOQ 30 mg in atopic dermatitis clinical trials. When notable differences in frequency were observed between indications, these are presented in the footnotes below the table.

**Table 23. Adverse reactions**

System Organ Class	Very common	Common	Uncommon
Infections and infestations	Upper respiratory tract infections (URTI) <sup>a</sup>	Bronchitis <sup>a,b</sup> Herpes zoster Herpes simplex <sup>a</sup> Folliculitis Influenza Urinary tract infection	Pneumonia Oral candidiasis Diverticulitis
Blood and lymphatic system disorders		Anaemia Neutropaenia <b>Lymphopaenia</b>	
Metabolism and nutrition disorders		Hypercholesterolaemia <sup>b</sup> <b>Hyperlipidaemia<sup>a,b</sup></b>	Hypertriglyceridaemia
Respiratory, thoracic and mediastinal disorders		Cough	
Gastrointestinal disorders		Abdominal pain <sup>a,d</sup> Nausea	
Skin and subcutaneous tissue disorders	Acne <sup>c,d</sup>	Urticaria <sup>c</sup> <b>Rash<sup>a</sup></b>	
General disorders and administration site conditions		Fatigue Pyrexia	
Investigations		Blood CPK increased ALT increased <sup>b</sup> AST increased <sup>b</sup> Weight increased	
Nervous system disorders		Headache	
<p><sup>a</sup> Presented as grouped term</p> <p><sup>b</sup> In atopic dermatitis trials, the frequency of bronchitis, hypercholesterolaemia, <b>hyperlipidaemia</b>, ALT increased, and AST increased was uncommon.</p> <p><sup>c</sup> In rheumatologic disease trials, the frequency was common for acne and uncommon for urticaria</p> <p><sup>d</sup> <b>In ulcerative colitis trials, the frequency was common for acne; abdominal pain was less frequent for upadacitinib than for placebo.</b></p>			

Description of selected adverse reactions

Rheumatoid arthritis

Infections

In placebo-controlled clinical studies with background DMARDs, the frequency of infection over 12/14 weeks in the upadacitinib 15 mg group was 27.4% compared to 20.9% in the placebo group. In methotrexate (MTX)-controlled studies, the frequency of infection over 12/14 weeks in the upadacitinib 15 mg monotherapy group was 19.5% compared to 24.0% in the MTX group. The overall long-term rate of infections for the upadacitinib 15 mg group across all five Phase 3 clinical studies (2,630 patients) was 93.7 events per 100 patient-years.

In placebo-controlled clinical studies with background DMARDs, the frequency of serious infection over 12/14 weeks in the upadacitinib 15 mg group was 1.2% compared to 0.6% in the placebo group. In MTX-controlled studies, the frequency of serious infection over 12/14 weeks in the upadacitinib 15 mg monotherapy group was 0.6% compared to 0.4% in the MTX group. The overall long-term rate

of serious infections for the upadacitinib 15 mg group across all five Phase 3 clinical studies was 3.8 events per 100 patient-years. The most common serious infection was pneumonia. The rate of serious infections remained stable with long-term exposure.

#### *Opportunistic infections (excluding tuberculosis)*

In placebo-controlled clinical studies with background DMARDs, the frequency of opportunistic infections over 12/14 weeks in the upadacitinib 15 mg group was 0.5% compared to 0.3% in the placebo group. In MTX-controlled studies, there were no cases of opportunistic infection over 12/14 weeks in the upadacitinib 15 mg monotherapy group and 0.2% in the MTX group. The overall long-term rate of opportunistic infections for the upadacitinib 15 mg group across all five Phase 3 clinical studies was 0.6 events per 100 patient-years.

The long-term rate of herpes zoster for the upadacitinib 15 mg group across all five Phase 3 clinical studies was 3.7 events per 100 patient-years. Most of the herpes zoster events involved a single dermatome and were non-serious.

#### *Hepatic transaminase elevations*

In placebo-controlled studies with background DMARDs, for up to 12/14 weeks, alanine transaminase (ALT) and aspartate transaminase (AST) elevations  $\geq 3$  x upper limit of normal (ULN) in at least one measurement were observed in 2.1% and 1.5% of patients treated with upadacitinib 15 mg, compared to 1.5% and 0.7%, respectively, of patients treated with placebo. Most cases of hepatic transaminase elevations were asymptomatic and transient.

In MTX-controlled studies, for up to 12/14 weeks, ALT and AST elevations  $\geq 3$  x ULN in at least one measurement were observed in 0.8% and 0.4% of patients treated with upadacitinib 15 mg, compared to 1.9% and 0.9%, respectively, of patients treated with MTX.

The pattern and incidence of elevation in ALT/AST remained stable over time including in ~~long~~-~~long~~-term extension studies.

#### *Lipid elevations*

Upadacitinib 15 mg treatment was associated with increases in lipid parameters including total cholesterol, triglycerides, LDL cholesterol and HDL cholesterol. There was no change in the LDL/HDL ratio. Elevations were observed at 2 to 4 weeks of treatment and remained stable with longer-term treatment. Among patients in the controlled studies with baseline values below the specified limits, the following frequencies of patients were observed to shift to above the specified limits on at least one occasion during 12/14 weeks (including patients who had an isolated elevated value):

- Total cholesterol  $\geq 5.17$  mmol/L (200 mg/dL): 62% vs. 31%, in the upadacitinib 15 mg and placebo groups, respectively
- LDL cholesterol  $\geq 3.36$  mmol/L (130 mg/dL): 42% vs. 19%, in the upadacitinib 15 mg and placebo groups, respectively
- HDL cholesterol  $\geq 1.03$  mmol/L (40 mg/dL): 89% vs. 61%, in the upadacitinib 15 mg and placebo groups, respectively
- Triglycerides  $\geq 2.26$  mmol/L (200 mg/dL): 25% vs. 15%, in the upadacitinib 15 mg and placebo groups, respectively

#### *Creatine phosphokinase*

In placebo-controlled studies with background DMARDs, for up to 12/14 weeks, increases in CPK values were observed. CPK elevations  $> 5$  x upper limit of normal (ULN) were reported in 1.0% and

0.3% of patients over 12/14 weeks in the upadacitinib 15 mg and placebo groups, respectively. Most elevations > 5 x ULN were transient and did not require treatment discontinuation. Mean CPK values increased by 4 weeks with a mean increase of 60 U/L at 12 weeks and then remained stable at an increased value thereafter including with extended therapy.

### *Neutropaenia*

In placebo-controlled studies with background DMARDs, for up to 12/14 weeks, decreases in neutrophil counts below  $1 \times 10^9$  cells/L in at least one measurement occurred in 1.1% and <0.1% of patients in the upadacitinib 15 mg and placebo groups, respectively. In clinical studies, treatment was interrupted in response to ANC <  $1 \times 10^9$  cells/L (see section 4.2). Mean neutrophil counts decreased over 4 to 8 weeks. The decreases in neutrophil counts remained stable at a lower value than baseline over time including with extended therapy.

### *Psoriatic arthritis*

Overall, the safety profile observed in patients with active psoriatic arthritis treated with upadacitinib 15 mg was consistent with the safety profile observed in patients with rheumatoid arthritis. A higher rate of serious infections (2.6 events per 100 patient-years and 1.3 events per 100 patient-years, respectively) and hepatic transaminase elevations (ALT elevations Grade 3 and higher rates 1.4% and 0.4%, respectively) was observed in patients treated with upadacitinib in combination with MTX therapy compared to patients treated with monotherapy.

### *Ankylosing spondylitis*

Overall, the safety profile observed in patients with active ankylosing spondylitis treated with upadacitinib 15 mg was consistent with the safety profile observed in patients with rheumatoid arthritis. No new safety findings were identified.

### *Atopic dermatitis*

#### *Infections*

In the placebo-controlled period of the clinical studies, the frequency of infection over 16 weeks in the upadacitinib 15 mg and 30 mg groups was 39% and 43% compared to 30% in the placebo group, respectively. The long-term rate of infections for the upadacitinib 15 mg and 30 mg groups was 98.5 and 109.6 events per 100 patient-years, respectively.

In placebo-controlled clinical studies, the frequency of serious infection over 16 weeks in the upadacitinib 15 mg and 30 mg groups was 0.8% and 0.4% compared to 0.6% in the placebo group, respectively. The long-term rate of serious infections for the upadacitinib 15 mg and 30 mg groups was 2.3 and 2.8 events per 100 patient-years, respectively.

#### *Opportunistic infections (excluding tuberculosis)*

In the placebo-controlled period of the clinical studies, all opportunistic infections (excluding TB and herpes zoster) reported were *eczema herpeticum*. The frequency of *eczema herpeticum* over 16 weeks in the upadacitinib 15 mg and 30 mg groups was 0.7% and 0.8% compared to 0.4% in the placebo group, respectively. The long-term rate of *eczema herpeticum* for the upadacitinib 15 mg and 30 mg groups was 1.6 and 1.8 events per 100 patient-years, respectively. One case of esophageal candidiasis was reported with upadacitinib 30 mg.

The long-term rate of herpes zoster for the upadacitinib 15 mg and 30 mg groups was 3.5 and 5.2 events per 100 patient-years, respectively. Most of the herpes zoster events involved a single dermatome and were non-serious.

### *Laboratory abnormalities*

Dose-dependent changes in ALT increased and/or AST increased ( $\geq 3 \times \text{ULN}$ ), lipid parameters, CPK values ( $> 5 \times \text{ULN}$ ), and neutropaenia ( $\text{ANC} < 1 \times 10^9 \text{ cells/L}$ ) associated with upadacitinib treatment were similar to what was observed in the rheumatologic disease clinical studies.

Small increases in LDL cholesterol were observed after week 16 in atopic dermatitis studies.

### *Ulcerative colitis*

The overall safety profile observed in patients with ulcerative colitis was generally consistent with that observed in patients with rheumatoid arthritis.

A higher rate of herpes zoster was observed with an induction treatment period of 16 weeks vs 8 weeks.

### *Infections*

In the placebo-controlled induction studies, the frequency of infection over 8 weeks in the upadacitinib 45 mg group compared to the placebo group was 20.7% and 17.5%, respectively. In the placebo-controlled maintenance study, the frequency of infection over 52 weeks in the upadacitinib 15 mg and 30 mg groups was 38.4% and 40.6%, respectively, compared to 37.6% in the placebo group. The long-term rate of infections for upadacitinib 15 mg and 30 mg was 73.8 and 82.6 events per 100 patient-years, respectively.

In the placebo-controlled induction studies, the frequency of serious infection over 8 weeks in both the upadacitinib 45 mg group and the placebo group was 1.3%. No additional serious infections were observed over 8-week extended treatment with upadacitinib 45 mg. In the placebo-controlled maintenance study, the frequency of serious infection over 52 weeks in the upadacitinib 15 mg and 30 mg groups was 3.2% and 2.4%, respectively, compared to 3.3% in the placebo group. The long-term rate of serious infections for the upadacitinib 15 mg and 30 mg groups was 4.1 and 3.9 events per 100 patient-years, respectively. The most frequently reported serious infection in the induction and maintenance phases was COVID-19 pneumonia.

### *Opportunistic infections (excluding tuberculosis)*

In the placebo-controlled induction studies over 8 weeks, the frequency of opportunistic infection (excluding tuberculosis and herpes zoster) in the upadacitinib 45 mg group was 0.4% and 0.3% in the placebo group. No additional opportunistic infections (excluding tuberculosis and herpes zoster) were observed over 8-week extended treatment with upadacitinib 45 mg. In the placebo-controlled maintenance study over 52 weeks, the frequency of opportunistic infection (excluding tuberculosis and herpes zoster) in the upadacitinib 15 mg and 30 mg groups was 0.8% and 0.4%, respectively, compared to 0.8% in the placebo group. The long-term rate of opportunistic infections (excluding tuberculosis and herpes zoster) for the upadacitinib 15 mg and 30 mg groups was 0.6 and 0.3 events per 100 patient-years, respectively.

In the placebo-controlled induction studies over 8 weeks, the frequency of herpes zoster in the upadacitinib 45 mg group was 0.6% and 0% in the placebo group. The frequency of herpes zoster was 3.9% over 16-week treatment with upadacitinib 45 mg. In the placebo-controlled maintenance study over 52 weeks, the frequency of herpes zoster in the upadacitinib 15 mg and 30 mg groups was 4.4% and 4.0%, respectively, compared to 0% in the placebo group. The long-term rate of herpes zoster for the upadacitinib 15 mg and 30 mg groups was 5.7 and 6.3 events per 100 patient-years, respectively.

### *Laboratory abnormalities*

In the induction and maintenance clinical studies, the laboratory changes in ALT increased and/or AST increased ( $\geq 3 \times \text{ULN}$ ), CPK values ( $> 5 \times \text{ULN}$ ), and neutropaenia ( $\text{ANC} < 1 \times 10^9 \text{ cells/L}$ ) associated with upadacitinib treatment were generally similar to what was observed in the



rheumatologic disease and atopic dermatitis clinical studies. Dose-dependent changes for these laboratory parameters associated with 15 mg and 30 mg upadacitinib treatment were observed:

In the placebo-controlled induction studies for up to 8 weeks, decreases in lymphocyte counts below  $0.5 \times 10^9$  cells/L in at least one measurement occurred in 2.0% and 0.8% of patients in the upadacitinib 45 mg and placebo groups, respectively. In the placebo-controlled maintenance study, for up to 52 weeks, decreases in lymphocyte counts below  $0.5 \times 10^9$  cells/L in at least one measurement occurred in 1.6%, 0.8% and 0.8% of patients in the upadacitinib 15 mg, 30 mg and placebo groups, respectively. In clinical studies, treatment was interrupted in response to  $ALC < 0.5 \times 10^9$  cells/L (see section 4.2). No notable mean changes of lymphocyte counts were observed during upadacitinib treatment over time.

Elevations in lipid parameters were observed at 8 weeks of treatment with upadacitinib 45 mg and remained generally stable with longer-term treatment with upadacitinib 15 mg and 30 mg. Among patients in the placebo-controlled induction studies with baseline values below the specified limits, the following frequencies of patients were observed to shift to above the specified limits on at least one occasion during 8 weeks (including patients who had an isolated elevated value):

- Total cholesterol  $\geq 5.17$  mmol/L (200 mg/dL): 49% vs. 11%, in the upadacitinib 45 mg and placebo groups, respectively
- LDL cholesterol  $\geq 3.36$  mmol/L (130 mg/dL): 27% vs. 9%, in the upadacitinib 45 mg and placebo groups, respectively
- HDL cholesterol  $\geq 1.03$  mmol/L (40 mg/dL): 79% vs. 36%, in the upadacitinib 45 mg and placebo groups, respectively
- Triglycerides  $\geq 2.26$  mmol/L (200 mg/dL): 6% vs 4% in the upadacitinib 45 mg and placebo groups, respectively

### Elderly

Based on limited data in atopic dermatitis patients aged 65 years and older, there was a higher rate of overall adverse reactions-with the upadacitinib 30 mg dose compared to the 15 mg dose.

Based on the limited data in ulcerative colitis patients aged 65 years and older, there was a higher rate of overall adverse reactions with the upadacitinib 30 mg dose compared to the 15 mg dose with maintenance treatment (see section 4.4).

### Paediatric population

A total of 343 adolescents aged 12 to 17 years with atopic dermatitis were treated in the Phase 3 studies, of whom 167 were exposed to 15 mg. The safety profile for upadacitinib 15 mg in adolescents was similar to that in adults. The safety and efficacy of the 30 mg dose in adolescents are still being investigated.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form:

<https://sideeffects.health.gov.il>

## **4.9 Overdose**



Upadacitinib was administered in clinical studies up to doses equivalent in daily AUC to 60 mg prolonged-release once daily. Adverse reactions were comparable to those seen at lower doses and no specific toxicities were identified. Approximately 90% of upadacitinib in the systemic circulation is eliminated within 24 hours of dosing (within the range of doses evaluated in clinical studies). In case of an overdose, it is recommended that the patient be monitored for signs and symptoms of adverse reactions. Patients who develop adverse reactions should receive appropriate treatment.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Immunosuppressants, selective immunosuppressants ATC code: L04AA44

#### Mechanism of action

Upadacitinib is a selective and reversible Janus kinase (JAK) inhibitor. JAKs are intracellular enzymes that transmit cytokine or growth factor signals involved in a broad range of cellular processes including inflammatory responses, hematopoiesis, and immune surveillance. The JAK family of enzymes contains four members, JAK1, JAK2, JAK3 and TYK2 which work in pairs to phosphorylate and activate signal transducers and activators of transcription (STATs). This phosphorylation, in turn, modulates gene expression and cellular function. JAK1 is important in inflammatory cytokine signals while JAK2 is important for red blood cell maturation and JAK3 signals play a role in immune surveillance and lymphocyte function.

In human cellular assays, upadacitinib preferentially inhibits signalling by JAK1 or JAK1/3 with functional selectivity over cytokine receptors that signal via pairs of JAK2. Atopic dermatitis is driven by pro-inflammatory cytokines (including IL-4, IL-13, IL-22, TSLP, IL-31 and IFN- $\gamma$ ) that transduce signals via the JAK1 pathway. Inhibiting JAK1 with upadacitinib reduces the signaling of many mediators which drive the signs and symptoms of atopic dermatitis such as eczematous skin lesions and pruritus. [Pro-inflammatory cytokines \(primarily IL-6, IL-7, IL-15 and IFN \$\gamma\$ \) transduce signals via the JAK1 pathway and are involved in ulcerative colitis pathogenesis. JAK1 inhibition with upadacitinib modulates the signalling of the JAK-dependent cytokines underlying the inflammatory burden and signs and symptoms of ulcerative colitis.](#)

#### Pharmacodynamic effects

##### *Inhibition of IL-6 induced STAT3 and IL-7 induced STAT5 phosphorylation*

In healthy volunteers, the administration of upadacitinib (immediate-release formulation) resulted in a dose- and concentration-dependent inhibition of IL-6 (JAK1/JAK2) - induced STAT3 and IL-7 (JAK1/JAK3)-induced STAT5 phosphorylation in whole blood. The maximal inhibition was observed 1 hour after dosing which returned to near baseline by the end of dosing interval.

##### *Lymphocytes*

In patients with rheumatoid arthritis, treatment with upadacitinib was associated with a small, transient increase in mean ALC from baseline up to week 36 which gradually returned to at or near baseline levels with continued treatment.

##### *hsCRP*

In patients with rheumatoid arthritis, treatment with upadacitinib was associated with decreases from baseline in mean hsCRP levels as early as week 1 which were maintained with continued treatment.

##### *Vaccine study*

The influence of upadacitinib on the humoral response following the administration of inactivated pneumococcal polysaccharide conjugate vaccine (13-valent, adsorbed) was evaluated in 111 patients with rheumatoid arthritis under stable treatment with upadacitinib 15 mg (n=87) or 30 mg (n=24). 97% of patients (n=108) were on concomitant methotrexate. The primary endpoint was the proportion of patients with satisfactory humoral response defined as  $\geq 2$ -fold increase in antibody concentration from baseline to week 4 in at least 6 out of the 12 pneumococcal antigens (1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F). Results at week 4 demonstrated a satisfactory humoral response in 67.5% (95% CI: 57.4, 77.5) and 56.5% (95% CI: 36.3, 76.8) of patients treated with upadacitinib 15 mg and 30 mg, respectively.

### Clinical efficacy and safety

#### *Rheumatoid arthritis*

The efficacy and safety of upadacitinib 15 mg once daily was assessed in five Phase 3 randomised, double-blind, multicentre studies in patients with moderately to severely active rheumatoid arthritis and fulfilling the ACR/EULAR 2010 classification criteria (see Table 3.4). Patients 18 years of age and older were eligible to participate. The presence of at least 6 tender and 6 swollen joints and evidence of systemic inflammation based on elevation of hsCRP was required at baseline. Four studies included long-term extensions for up to 5 years, and one study (SELECT-COMPARE) included a long-term extension for up to 10 years.

The primary analysis for each of these studies included all randomised subjects who received at least 1 dose of upadacitinib or placebo, and non-responder imputation was used for categorical endpoints.

Across the Phase 3 studies, the efficacy seen with upadacitinib 15 mg QD was generally similar to that observed with upadacitinib 30 mg QD.

**Table 3.4 Clinical trials summary**

<b>Study name</b>	<b>Population (n)</b>	<b>Treatment arms</b>	<b>Key outcome measures</b>
SELECT-EARLY	MTX-naïve <sup>a</sup> (947)	<ul style="list-style-type: none"> <li>• Upadacitinib 15 mg</li> <li>• Upadacitinib 30 mg</li> <li>• MTX</li> </ul> <p>Monotherapy</p>	<ul style="list-style-type: none"> <li>• Primary endpoint: clinical remission (DAS28-CRP) at week 24</li> <li>• Low disease activity (DAS28-CRP)</li> <li>• ACR50</li> <li>• Radiographic progression (mTSS)</li> <li>• Physical function (HAQ-DI)</li> <li>• SF-36 PCS</li> </ul>
SELECT-MONOTHERAPY	MTX-IR <sup>b</sup> (648)	<ul style="list-style-type: none"> <li>• Upadacitinib 15 mg</li> <li>• Upadacitinib 30 mg</li> <li>• MTX</li> </ul> <p>Monotherapy</p>	<ul style="list-style-type: none"> <li>• Primary endpoint: low disease activity (DAS28-CRP) at week 14</li> <li>• Clinical remission (DAS28-CRP)</li> <li>• ACR20</li> <li>• Physical function (HAQ-DI)</li> <li>• SF-36 PCS</li> <li>• Morning stiffness</li> </ul>
SELECT-NEXT	csDMARD-IR <sup>c</sup> (661)	<ul style="list-style-type: none"> <li>• Upadacitinib 15 mg</li> <li>• Upadacitinib 30 mg</li> <li>• Placebo</li> </ul> <p>On background csDMARDs</p>	<ul style="list-style-type: none"> <li>• Primary endpoint: low disease activity (DAS28-CRP) at week 12</li> <li>• Clinical remission (DAS28-CRP)</li> <li>• ACR20</li> <li>• Physical function (HAQ-DI)</li> <li>• SF-36 PCS</li> <li>• Low disease activity (CDAI)</li> <li>• Morning stiffness</li> <li>• FACIT-F</li> </ul>

SELECT-COMPARE	MTX-IR <sup>d</sup> (1,629)	<ul style="list-style-type: none"> <li>• Upadacitinib 15 mg</li> <li>• Placebo</li> <li>• Adalimumab 40 mg</li> </ul> <p>On background MTX</p>	<ul style="list-style-type: none"> <li>• Primary endpoint: clinical remission (DAS28-CRP) at week 12</li> <li>• Low disease activity (DAS28-CRP)</li> <li>• ACR20</li> <li>• Low disease activity (DAS28-CRP) vs adalimumab</li> <li>• Radiographic progression (mTSS)</li> <li>• Physical function (HAQ-DI)</li> <li>• SF-36 PCS</li> <li>• Low disease activity (CDAI)</li> <li>• Morning stiffness</li> <li>• FACIT-F</li> </ul>
SELECT-BEYOND	bDMARD-IR <sup>e</sup> (499)	<ul style="list-style-type: none"> <li>• Upadacitinib 15 mg</li> <li>• Upadacitinib 30 mg</li> <li>• Placebo</li> </ul> <p>On background csDMARDs</p>	<ul style="list-style-type: none"> <li>• Primary endpoint: low disease activity (DAS28-CRP) at week 12</li> <li>• ACR20</li> <li>• Physical function (HAQ-DI)</li> <li>• SF-36 PCS</li> </ul>

Abbreviations: ACR20 (or 50) = American College of Rheumatology  $\geq 20\%$  (or  $\geq 50\%$ ) improvement; bDMARD = biologic disease-modifying anti-rheumatic drug, CRP = C-Reactive Protein, DAS28 = Disease Activity Score 28 joints, mTSS = modified Total Sharp Score, csDMARD = conventional synthetic disease-modifying anti-rheumatic drug, HAQ-DI = Health Assessment Questionnaire Disability Index, SF-36 PCS = Short Form (36) Health Survey (SF-36) Physical Component Summary, CDAI = Clinical Disease Activity Index, FACIT-F = Functional Assessment of Chronic Illness Therapy-Fatigue score, IR = inadequate responder, MTX = methotrexate, n = number randomised

<sup>a</sup> Patients were naïve to MTX or received no more than 3 weekly MTX doses

<sup>b</sup> Patients had inadequate response to MTX

<sup>c</sup> Patients who had an inadequate response to csDMARDs; patients with prior exposure to at most one bDMARD were eligible (up to 20% of total number of patients) if they had either limited exposure (<3 months) or had to discontinue the bDMARD due to intolerability

<sup>d</sup> Patients who had an inadequate response to MTX; patients with prior exposure to at most one bDMARD (except adalimumab) were eligible (up to 20% of total study number of patients) if they had either limited exposure (<3 months) or had to discontinue the bDMARD due to intolerability

<sup>e</sup> Patients who had an inadequate response or intolerance to at least one bDMARD

### Clinical response

#### Remission and low disease activity

In the studies, a significantly higher proportion of patients treated with upadacitinib 15 mg achieved low disease activity (DAS28-CRP  $\leq 3.2$ ) and clinical remission (DAS28-CRP  $< 2.6$ ) compared to placebo, MTX or adalimumab (Table 45). Compared to adalimumab, significantly higher rates of low disease activity were achieved at week 12 in SELECT-COMPARE. Overall, both low disease activity and clinical remission rates were consistent across patient populations, with or without MTX. At 3 years, 297/651 (45.6%) and 111/327 (33.9%) patients remained on originally randomised treatment of upadacitinib 15 mg or adalimumab, respectively, in SELECT-COMPARE, and 216/317 (68.1%) and 149/315 (47.3%) patients remained on originally randomised treatment of upadacitinib 15 mg or MTX monotherapy, respectively, in SELECT-EARLY. Among the patients who remained on their originally allocated treatment, low disease activity and clinical remission were maintained through 3 years.

#### ACR response

In all studies, more patients treated with upadacitinib 15 mg achieved ACR20, ACR50, and ACR70 responses at 12 weeks compared to placebo, MTX, or adalimumab (Table 45). Time to onset of

efficacy was rapid across measures with greater responses seen as early as week 1 for ACR20. Durable response rates were observed (with or without MTX), with ACR20/50/70 responses maintained through 3 years among the patients who remained on their originally allocated treatment.

Treatment with upadacitinib 15 mg, alone or in combination with csDMARDs, resulted in improvements in individual ACR components, including tender and swollen joint counts, patient and physician global assessments, HAQ-DI, pain assessment and hsCRP.

**Table 45: Response and remission**

Study	SELECT EARLY MTX-Naïve		SELECT MONO MTX-IR		SELECT NEXT csDMARD-IR		SELECT COMPARE MTX-IR			SELECT BEYOND bDMARD-IR	
	MTX	UPA 15mg	MTX	UPA 15mg	PBO	UPA 15mg	PBO	UPA 15mg	ADA 40mg	PBO	UPA 15mg
N	314	317	216	217	221	221	651	651	327	169	164
Week											
<b>LDA DAS28-CRP ≤3.2 (% of patients)</b>											
12 <sup>a</sup> /14 <sup>b</sup>	28	53 <sup>g</sup>	19	45 <sup>e</sup>	17	48 <sup>e</sup>	14	45 <sup>e,h</sup>	29	14	43 <sup>e</sup>
24 <sup>c</sup> /26 <sup>d</sup>	32	60 <sup>f</sup>					18	55 <sup>g,h</sup>	39		
48	39	59 <sup>g</sup>						50 <sup>h</sup>	35		
<b>CR DAS28-CRP &lt;2.6 (% of patients)</b>											
12 <sup>a</sup> /14 <sup>b</sup>	14	36 <sup>g</sup>	8	28 <sup>e</sup>	10	31 <sup>e</sup>	6	29 <sup>e,h</sup>	18	9	29 <sup>g</sup>
24 <sup>c</sup> /26 <sup>d</sup>	18	48 <sup>e</sup>					9	41 <sup>g,h</sup>	27		
48	29	49 <sup>g</sup>						38 <sup>i</sup>	28		
<b>ACR20 (% of patients)</b>											
12 <sup>a</sup> /14 <sup>b</sup>	54	76 <sup>g</sup>	41	68 <sup>e</sup>	36	64 <sup>e</sup>	36	71 <sup>e,j</sup>	63	28	65 <sup>e</sup>
24 <sup>c</sup> /26 <sup>d</sup>	59	79 <sup>g</sup>					36	67 <sup>g,i</sup>	57		
48	57	74 <sup>g</sup>						65 <sup>i</sup>	54		
<b>ACR50 (% of patients)</b>											
12 <sup>a</sup> /14 <sup>b</sup>	28	52 <sup>g</sup>	15	42 <sup>g</sup>	15	38 <sup>g</sup>	15	45 <sup>g,h</sup>	29	12	34 <sup>g</sup>
24 <sup>c</sup> /26 <sup>d</sup>	33	60 <sup>e</sup>					21	54 <sup>g,h</sup>	42		
48	43	63 <sup>g</sup>						49 <sup>i</sup>	40		
<b>ACR70 (% of patients)</b>											
12 <sup>a</sup> /14 <sup>b</sup>	14	32 <sup>g</sup>	3	23 <sup>g</sup>	6	21 <sup>g</sup>	5	25 <sup>g,h</sup>	13	7	12
24 <sup>c</sup> /26 <sup>d</sup>	18	44 <sup>g</sup>					10	35 <sup>g,h</sup>	23		
48	29	51 <sup>g</sup>						36 <sup>h</sup>	23		
<b>CDAI ≤10 (% of patients)</b>											
12 <sup>a</sup> /14 <sup>b</sup>	30	46 <sup>g</sup>	25	35 <sup>l</sup>	19	40 <sup>e</sup>	16	40 <sup>e,h</sup>	30	14	32 <sup>g</sup>
24 <sup>c</sup> /26 <sup>d</sup>	38	56 <sup>g</sup>					22	53 <sup>g,h</sup>	38		
48	43	60 <sup>g</sup>						47 <sup>h</sup>	34		
Abbreviations: ACR20 (or 50 or 70) = American College of Rheumatology ≥20% (or ≥50% or ≥70%) improvement; ADA = adalimumab; CDAI = Clinical Disease Activity Index; CR = Clinical Remission; CRP = C-Reactive Protein, DAS28 = Disease Activity Score 28 joints; IR = inadequate responder; LDA = Low Disease Activity; MTX = methotrexate; PBO = placebo; UPA= upadacitinib <sup>a</sup> SELECT-NEXT, SELECT-EARLY, SELECT-COMPARE, SELECT-BEYOND <sup>b</sup> SELECT-MONOTHERAPY <sup>c</sup> SELECT-EARLY <sup>d</sup> SELECT-COMPARE <sup>e</sup> multiplicity-controlled p≤0.001 upadacitinib vs placebo or MTX comparison <sup>f</sup> multiplicity-controlled p≤0.01 upadacitinib vs placebo or MTX comparison <sup>g</sup> nominal p≤0.001 upadacitinib vs placebo or MTX comparison											

<sup>h</sup> nominal  $p \leq 0.001$  upadacitinib vs adalimumab comparison  
<sup>i</sup> nominal  $p \leq 0.01$  upadacitinib vs adalimumab comparison  
<sup>j</sup> nominal  $p < 0.05$  upadacitinib vs adalimumab comparison  
<sup>k</sup> nominal  $p \leq 0.01$  upadacitinib vs placebo or MTX comparison  
<sup>l</sup> nominal  $p < 0.05$  upadacitinib vs MTX comparison  
Note: Week 48-data derived from analysis on Full Analysis set (FAS) by randomised group using Non-Responder Imputation

### Radiographic response

Inhibition of progression of structural joint damage was assessed using the modified Total Sharp Score (mTSS) and its components, the erosion score and joint space narrowing score, at weeks 24/26 and week 48 in SELECT-EARLY and SELECT-COMPARE.

Treatment with upadacitinib 15 mg resulted in significantly greater inhibition of the progression of structural joint damage compared to placebo in combination with MTX in SELECT-COMPARE and as monotherapy compared to MTX in SELECT-EARLY (Table 56). Analyses of erosion and joint space narrowing scores were consistent with the overall scores. The proportion of patients with no radiographic progression (mTSS change  $\leq 0$ ) was significantly higher with upadacitinib 15 mg in both studies. Inhibition of progression of structural joint damage was maintained through week 96 in both studies for patients who remained on their originally allocated treatment with upadacitinib 15 mg (based on available results from 327 patients in SELECT-COMPARE and 238 patients in SELECT-EARLY).

**Table 56: Radiographic changes**

Study	SELECT EARLY MTX-Naïve		SELECT COMPARE MTX-IR		
	MTX	UPA 15 mg	PBO <sup>a</sup>	UPA 15 mg	ADA 40 mg
<b>Modified Total Sharp Score, mean change from baseline</b>					
Week 24 <sup>b</sup> /26 <sup>c</sup>	0.7	0.1 <sup>f</sup>	0.9	0.2 <sup>g</sup>	0.1
Week 48	1.0	0.03 <sup>e</sup>	1.7	0.3 <sup>e</sup>	0.4
<b>Proportion of patients with no radiographic progression<sup>d</sup></b>					
Week 24 <sup>b</sup> /26 <sup>c</sup>	77.7	87.5 <sup>f</sup>	76.0	83.5 <sup>f</sup>	86.8
Week 48	74.3	89.9 <sup>e</sup>	74.1	86.4 <sup>e</sup>	87.9
Abbreviations: ADA = adalimumab; IR = inadequate responder; MTX = methotrexate; PBO = placebo; UPA= upadacitinib					
<sup>a</sup> All placebo data at week 48 derived using linear extrapolation					
<sup>b</sup> SELECT-EARLY					
<sup>c</sup> SELECT-COMPARE					
<sup>d</sup> No progression defined as mTSS change $\leq 0$					
<sup>e</sup> nominal $p \leq 0.001$ upadacitinib vs placebo or MTX comparison					
<sup>f</sup> multiplicity-controlled $p \leq 0.01$ upadacitinib vs placebo or MTX comparison					
<sup>g</sup> multiplicity-controlled $p \leq 0.001$ upadacitinib vs placebo or MTX comparison					

### Physical function response and health-related outcomes

Treatment with upadacitinib 15 mg, alone or in combination with csDMARDs, resulted in a significantly greater improvement in physical function compared to all comparators as measured by HAQ-DI (see Table 67). Improvement in HAQ-DI was maintained through 3 years for patients who remained on their originally allocated treatment with upadacitinib 15 mg based on available results from SELECT-COMPARE and SELECT-EARLY.

**Table 67: Mean change from baseline in HAQ-DI<sup>a,b</sup>**

Study	SELECT EARLY MTX-Naïve		SELECT MONO MTX-IR		SELECT NEXT csDMARD-IR		SELECT COMPARE MTX-IR			SELECT BEYOND BIO-IR	
	MTX	UPA 15mg	MTX	UPA 15mg	PBO	UPA 15mg	PBO	UPA 15mg	ADA 40mg	PBO	UPA 15mg
N	313	317	216	216	220	216	648	644	324	165	163
Baseline score, mean	1.6	1.6	1.5	1.5	1.4	1.5	1.6	1.6	1.6	1.6	1.7
Week 12 <sup>c</sup> /14 <sup>d</sup>	-0.5	-0.8 <sup>h</sup>	-0.3	-0.7 <sup>g</sup>	-0.3	-0.6 <sup>g</sup>	-0.3	-0.6 <sup>g,i</sup>	-0.5	-0.2	-0.4 <sup>g</sup>
Week 24 <sup>e</sup> /26 <sup>f</sup>	-0.6	-0.9 <sup>g</sup>					-0.3	-0.7 <sup>h,i</sup>	-0.6		

Abbreviations: ADA = adalimumab; HAQ-DI = Health Assessment Questionnaire-Disability Index; IR = inadequate responder; MTX = methotrexate; PBO = placebo; UPA = upadacitinib

<sup>a</sup> Data shown are mean

<sup>b</sup> Health Assessment Questionnaire-Disability Index: 0=best, 3=worst; 20 questions; 8 categories: dressing and grooming, arising, eating, walking, hygiene, reach, grip, and activities.

<sup>c</sup> SELECT-EARLY, SELECT-NEXT, SELECT-COMPARE, SELECT-BEYOND

<sup>d</sup> SELECT-MONOTHERAPY

<sup>e</sup> SELECT-EARLY

<sup>f</sup> SELECT-COMPARE

<sup>g</sup> multiplicity-controlled p≤0.001 upadacitinib vs placebo or MTX comparison

<sup>h</sup> nominal p≤0.001 upadacitinib vs placebo or MTX comparison

<sup>i</sup> nominal p≤0.01 upadacitinib vs adalimumab comparison

In the studies SELECT-MONOTHERAPY, SELECT-NEXT, and SELECT-COMPARE, treatment with upadacitinib 15 mg resulted in a significantly greater improvement in the mean duration of morning joint stiffness compared to placebo or MTX.

In the clinical studies, upadacitinib treated patients reported significant improvements in patient-reported quality of life, as measured by the Short Form (36) Health Survey (SF-36) Physical Component Summary compared to placebo and MTX. Moreover, upadacitinib treated patients reported significant improvements in fatigue, as measured by the Functional Assessment of Chronic Illness Therapy-Fatigue score (FACIT-F) compared to placebo.

### *Psoriatic arthritis*

The efficacy and safety of upadacitinib 15 mg once daily were assessed in two Phase 3 randomised, double-blind, multicentre, placebo-controlled studies in patients 18 years of age or older with moderately to severely active psoriatic arthritis. All patients had active psoriatic arthritis for at least 6 months based upon the Classification Criteria for Psoriatic Arthritis (CASPAR), at least 3 tender joints and at least 3 swollen joints, and active plaque psoriasis or history of plaque psoriasis. For both studies, the primary endpoint was the proportion of patients who achieved an ACR20 response at week 12.

SELECT-PsA 1 was a 24-week trial in 1705 patients who had an inadequate response or intolerance to at least one non-biologic DMARD. At baseline, 1393 (82%) of patients were on at least one concomitant non-biologic DMARD; 1084 (64%) of patients received concomitant MTX only; and 311 (18%) of patients were on monotherapy. Patients received upadacitinib 15 mg or 30 mg once daily, adalimumab, or placebo. At week 24, all patients randomised to placebo were switched to upadacitinib 15 mg or 30 mg once daily in a blinded manner. SELECT-PsA 1 included a long-term extension for up to 5 years.



SELECT-PsA 2 was a 24-week trial in 642 patients who had an inadequate response or intolerance to at least one biologic DMARD. At baseline, 296 (46%) of patients were on at least one concomitant non-biologic DMARD; 222 (35%) of patients received concomitant MTX only; and 345 (54%) of patients were on monotherapy. Patients received upadacitinib 15 mg or 30 mg once daily or placebo. At week 24, all patients randomised to placebo were switched to upadacitinib 15 mg or 30 mg once daily in a blinded manner. SELECT-PsA 2 included a long-term extension for up to 3 years.

### Clinical response

In both studies, a statistically significant greater proportion of patients treated with upadacitinib 15 mg achieved ACR20 response compared to placebo at week 12 (Table 7:8). Time to onset of efficacy was rapid across measures with greater responses seen as early as week 2 for ACR20.

Treatment with upadacitinib 15 mg resulted in improvements in individual ACR components, including tender/painful and swollen joint counts, patient and physician global assessments, HAQ-DI, pain assessment, and hsCRP compared to placebo.

In SELECT-PsA 1, upadacitinib 15 mg achieved non-inferiority compared to adalimumab in the proportion of patients achieving ACR20 response at week 12; however, superiority to adalimumab could not be demonstrated.

In both studies, consistent responses were observed alone or in combination with methotrexate for primary and key secondary endpoints.

The efficacy of upadacitinib 15 mg was demonstrated regardless of subgroups evaluated including baseline BMI, baseline hsCRP, and number of prior non-biologic DMARDs ( $\leq 1$  or  $>1$ ).

**Table 7:8 Clinical response in SELECT-PsA 1 and SELECT-PsA 2**

Study	SELECT-PsA 1 non-biologic DMARD-IR			SELECT-PsA 2 bDMARD-IR	
	PBO	UPA 15 mg	ADA 40 mg	PBO	UPA 15 mg
N	423	429	429	212	211
<b>ACR20, % of patients (95% CI)</b>					
Week 12	36 (32, 41)	71 (66, 75) <sup>f</sup>	65 (61, 70)	24 (18, 30)	57 (50, 64)
Difference from placebo (95% CI)	35 (28, 41) <sup>d,e</sup>		-	33 (24, 42) <sup>d,e</sup>	
Week 24	45 (40, 50)	73 (69, 78)	67 (63, 72)	20 (15, 26)	59 (53, 66)
Week 56		74 (70, 79)	69 (64, 73)		60 (53, 66)
<b>ACR50, % of patients (95% CI)</b>					
Week 12	13 (10, 17)	38 (33, 42)	38 (33, 42)	5 (2, 8)	32 (26, 38)
Week 24	19 (15, 23)	52 (48, 57)	44 (40, 49)	9 (6, 13)	38 (32, 45)
Week 56		60 (55, 64)	51 (47, 56)		41 (34, 47)
<b>ACR70, % of patients (95% CI)</b>					
Week 12	2 (1, 4)	16 (12, 19)	14 (11, 17)	1 (0, 1)	9 (5, 12)
Week 24	5 (3, 7)	29 (24, 33)	23 (19, 27)	1 (0, 2)	19 (14, 25)
Week 56		41 (36, 45)	31 (27, 36)		24 (18, 30)
<b>MDA, % of patients (95% CI)</b>					
Week 12	6 (4, 9)	25 (21, 29)	25 (21, 29)	4 (2, 7)	17 (12, 22)
Week 24	12 (9, 15)	37 (32, 41) <sup>e</sup>	33 (29, 38)	3 (1, 5)	25 (19, 31) <sup>e</sup>
Week 56		45 (40, 50)	40 (35, 44)		29 (23, 36)
<b>Resolution of enthesitis (LEI=0), % of patients (95% CI)<sup>a</sup></b>					

Week 12	33 (27, 39)	47 (42, 53)	47 (41, 53)	20 (14, 27)	39 (31, 47)
Week 24	32 (27, 39)	54 (48, 60) <sup>e</sup>	47 (42, 53)	15 (9, 21)	43 (34, 51)
Week 56		59 (53, 65)	54 (48, 60)		43 (34, 51)
<b>Resolution of dactylitis (LDI=0), % of patients (95% CI)<sup>b</sup></b>					
Week 12	42 (33, 51)	74 (66, 81)	72 (64, 80)	36 (24, 48)	64 (51, 76)
Week 24	40 (31, 48)	77 (69, 84)	74 (66, 82)	28 (17, 39)	58 (45, 71)
Week 56		75 (68, 82)	74 (66, 82)		51 (38, 64)
<b>PASI75, % of patients (95% CI)<sup>c</sup></b>					
Week 16	21 (16, 27)	63 (56, 69) <sup>e</sup>	53 (46, 60)	16 (10, 22)	52 (44, 61) <sup>e</sup>
Week 24	27 (21, 33)	64 (58, 70)	59 (52, 65)	19 (12, 26)	54 (45, 62)
Week 56		65 (59, 72)	61 (55, 68)		52 (44, 61)
<b>PASI90, % of patients (95% CI)<sup>c</sup></b>					
Week 16	12 (8, 17)	38 (32, 45)	39 (32, 45)	8 (4, 13)	35 (26, 43)
Week 24	17 (12, 22)	42 (35, 48)	45 (38, 52)	7 (3, 11)	36 (28, 44)
Week 56		49 (42, 56)	47 (40, 54)		41 (32, 49)
<p>Abbreviations: ACR20 (or 50 or 70) = American College of Rheumatology <math>\geq 20\%</math> (or <math>\geq 50\%</math> or <math>\geq 70\%</math>) improvement, ADA = adalimumab; bDMARD = biologic disease-modifying anti-rheumatic drug; IR = inadequate responder; MDA = minimal disease activity; PASI75 (or 90) = <math>\geq 75\%</math> (or <math>\geq 90\%</math>) improvement in Psoriasis Area and Severity Index; PBO = placebo; UPA= upadacitinib</p> <p>Patients who discontinued randomised treatment or were missing data at week of evaluation were imputed as non-responders in the analyses. For MDA, resolution of enthesitis, and resolution of dactylitis at week 24/56, the subjects rescued at week 16 were imputed as non-responders in the analyses.</p> <p><sup>a</sup> In patients with enthesitis at baseline (n=241, 270, and 265, respectively, for SELECT-PsA 1 and n=144 and 133, respectively, for SELECT-PsA 2)</p> <p><sup>b</sup> In patients with dactylitis at baseline (n=126, 136, and 127, respectively, for SELECT-PsA 1 and n=64 and 55, respectively, for SELECT-PsA 2)</p> <p><sup>c</sup> In patients with <math>\geq 3\%</math> BSA psoriasis at baseline (n=211, 214, and 211, respectively, for SELECT-PsA 1 and n=131 and 130, respectively, for SELECT-PsA 2)</p> <p><sup>d</sup> primary endpoint</p> <p><sup>e</sup> multiplicity-controlled <math>p \leq 0.001</math> upadacitinib vs placebo comparison</p> <p><sup>f</sup> multiplicity-controlled <math>p \leq 0.001</math> upadacitinib vs adalimumab comparison (non-inferiority test)</p>					

### Radiographic response

In SELECT-PsA 1, inhibition of progression of structural damage was assessed radiographically and expressed as the change from baseline in modified Total Sharp Score (mTSS) and its components, the erosion score and the joint space narrowing score, at week 24.

Treatment with upadacitinib 15 mg resulted in statistically significant greater inhibition of the progression of structural joint damage compared to placebo at week 24 (Table 89). Erosion and joint space narrowing scores were consistent with the overall scores. The proportion of patients with no radiographic progression (mTSS change  $\leq 0.5$ ) was higher with upadacitinib 15 mg compared to placebo at week 24.



**Table 98: Radiographic changes in SELECT-PsA 1**

Treatment Group	PBO	UPA 15 mg	ADA 40 mg
<b>Modified Total Sharp Score, mean change from baseline (95% CI)</b>			
Week 24	0.25 (0.13, 0.36)	-0.04 (-0.16, 0.07) <sup>c</sup>	0.01 (-0.11, 0.13)
Week 56 <sup>a</sup>	0.44 (0.29, 0.59)	-0.05 (-0.20, 0.09)	-0.06 (-0.20, 0.09)
<b>Proportion of patients with no radiographic progression<sup>b</sup>, % (95% CI)</b>			
Week 24	92 (89, 95)	96 (94, 98)	95 (93, 97)
Week 56 <sup>a</sup>	89 (86, 92)	97 (96, 99)	94 (92, 97)
Abbreviations: ADA = adalimumab; PBO = placebo; UPA= upadacitinib			
<sup>a</sup> All placebo data at week 56 derived using linear extrapolation			
<sup>b</sup> No progression defined as mTSS change ≤0.5			
<sup>c</sup> multiplicity-controlled p≤0.001 upadacitinib vs placebo comparison			

*Physical function response and health-related outcomes*

In SELECT-PsA 1, patients treated with upadacitinib 15 mg showed statistically significant improvement from baseline in physical function as assessed by HAQ-DI at week 12 (-0.42 [95% CI: -0.47, -0.37]) compared to placebo (-0.14 [95% CI: -0.18, -0.09]); improvement in patients treated with adalimumab was -0.34 (95% CI: -0.38, -0.29). In SELECT-PsA 2, patients treated with upadacitinib 15 mg showed statistically significant improvement from baseline in HAQDI at week 12 (-0.30 [95% CI: -0.37, -0.24]) compared to placebo (-0.10 [95% CI: -0.16, -0.03]). Improvement in physical function was maintained through week 56 in both studies.

Health-related quality of life was assessed by SF-36v2. In both studies, patients receiving upadacitinib 15 mg experienced statistically significant greater improvement from baseline in the Physical Component Summary score compared to placebo at week 12. Improvements from baseline were maintained through week 56 in both studies.

Patients receiving upadacitinib 15 mg experienced statistically significant improvement from baseline in fatigue, as measured by FACIT-F score, at week 12 compared to placebo in both studies. Improvements from baseline were maintained through week 56 in both studies.

At baseline, psoriatic spondylitis was reported in 31% and 34% of patients in SELECT-PsA 1 and SELECT-PsA 2, respectively. Patients with psoriatic spondylitis treated with upadacitinib 15 mg showed improvements from baseline in Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) scores compared to placebo at week 24. Improvements from baseline were maintained through week 56 in both studies.

*Ankylosing spondylitis*

The efficacy and safety of upadacitinib 15 mg once daily were assessed in two randomised, double-blind, multicentre, placebo-controlled studies in patients 18 years of age or older with active ankylosing spondylitis based upon the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) ≥4 and Patient's Assessment of Total Back Pain score ≥4. Both studies included long-term extensions for up to 2 years.

SELECT-AXIS 1 was a 14-week placebo-controlled trial in 187 ankylosing spondylitis patients with an inadequate response to at least two Nonsteroidal Anti-inflammatory Drugs (NSAIDs) or intolerance to or contraindication for NSAIDs and had no previous exposure to biologic DMARDs. At baseline, patients had symptoms of ankylosing spondylitis for an average of 14.4 years and approximately 16% of the patients were on a concomitant csDMARD. Patients received upadacitinib 15 mg once daily or placebo. At week 14, all patients randomised to placebo were switched to upadacitinib 15 mg once

daily. The primary endpoint was the proportion of patients achieving an Assessment of SpondyloArthritis international Society 40 (ASAS40) response at week 14.

SELECT-AXIS 2 (AS) was a 14-week placebo-controlled trial in 420 ankylosing spondylitis patients with prior exposure to bDMARDs (77.4% had lack of efficacy to either a tumour necrosis factor (TNF) blocker or interleukin-17 inhibitor (IL-17i); 30.2% had intolerance; 12.9% had prior exposure but not lack of efficacy to two bDMARDs). At baseline, patients had symptoms of ankylosing spondylitis for an average of 12.8 years and approximately 31% of the patients were on a concomitant csDMARD. Patients received upadacitinib 15 mg once daily or placebo. At week 14, all patients randomised to placebo were switched to upadacitinib 15 mg once daily. The primary endpoint was the proportion of patients achieving an Assessment of SpondyloArthritis international Society 40 (ASAS40) response at week 14.

### Clinical response

In both studies, a significantly greater proportion of patients treated with upadacitinib 15 mg achieved an ASAS40 response compared to placebo at week 14 (Table 9:10). A numerical difference between treatment groups was observed from week 2 in SELECT-AXIS 1 and week 4 in SELECT-AXIS 2 (AS) for ASAS40.

Treatment with upadacitinib 15 mg resulted in improvements in individual ASAS components (patient global assessment of disease activity, total back pain assessment, inflammation, and function) and other measures of disease activity, including hsCRP, at week 14 compared to placebo.

The efficacy of upadacitinib 15 mg was demonstrated regardless of subgroups evaluated including gender, baseline BMI, symptom duration of AS, baseline hsCRP, and prior use of bDMARDs.

**Table 9:10 Clinical response**

Study	SELECT-AXIS 1 bDMARD-naïve		SELECT-AXIS 2 (AS) bDMARD-IR	
	PBO	UPA 15 mg	PBO	UPA 15 mg
N	94	93	209	211
<b>ASAS40, % of patients (95% CI)<sup>a,b</sup></b>				
Week 14	25.5 (16.7, 34.3)	51.6 (41.5, 61.8)	18.2 (13.0, 23.4)	44.5 (37.8, 51.3)
Difference from placebo (95% CI)	26.1 (12.6, 39.5) <sup>c</sup>		26.4 (17.9, 34.9) <sup>c</sup>	
<b>ASAS20, % of patients (95% CI)<sup>a</sup></b>				
Week 14	40.4 (30.5, 50.3)	64.5 (54.8, 74.2) <sup>e</sup>	38.3 (31.7, 44.9)	65.4 (59.0, 71.8) <sup>c</sup>
<b>ASAS Partial Remission, % of patients (95% CI)</b>				
Week 14	1.1 (0.0, 3.1)	19.4 (11.3, 27.4) <sup>c</sup>	4.3 (1.6, 7.1)	17.5 (12.4, 22.7) <sup>c</sup>
<b>BASDAI 50, % of patients (95% CI)</b>				
Week 14	23.4 (14.8, 32.0)	45.2 (35.0, 55.3) <sup>d</sup>	16.7 (11.7, 21.8)	43.1 (36.4, 49.8) <sup>c</sup>
<b>Change from baseline in ASDAS-CRP (95% CI)</b>				
Week 14	-0.54 (-0.71, -0.37)	-1.45 (-1.62, -1.28) <sup>c</sup>	-0.49 (-0.62, -0.37)	-1.52 (-1.64, -1.39) <sup>c</sup>
<b>ASDAS Inactive Disease, % of patients (95% CI)</b>				
Week 14	0	16.1 (8.7, 23.6) <sup>c</sup>	1.9 (0.1, 3.8)	12.8 (8.3, 17.3) <sup>c</sup>

<b>ASDAS Low Disease Activity, % of patients (95% CI)</b>				
Week 14	10.6 (4.4, 16.9)	49.5 (39.3, 59.6) <sup>f</sup>	10.1 (6.0, 14.2)	44.1 (37.4, 50.8) <sup>e</sup>
<b>ASDAS Major Improvement, % of patients (95% CI)</b>				
Week 14	5.3 (0.8, 9.9)	32.3 (22.8, 41.8) <sup>e</sup>	4.8 (1.9, 7.7)	30.3 (24.1, 36.5) <sup>e</sup>

Abbreviations: ASAS20 (or ASAS40) = Assessment of SpondyloArthritis international Society  $\geq 20\%$  (or  $\geq 40\%$ ) improvement; ASDAS-CRP = Ankylosing Spondylitis Disease Activity Score C-Reactive Protein; BASDAI = Bath Ankylosing Spondylitis Disease Activity Index; PBO = placebo; UPA= upadacitinib

<sup>a</sup> An ASAS20 (ASAS40) response is defined as a  $\geq 20\%$  ( $\geq 40\%$ ) improvement and an absolute improvement from baseline of  $\geq 1$  ( $\geq 2$ ) unit(s) (range 0 to 10) in  $\geq 3$  of 4 domains (Patient Global, Total Back Pain, Function, and Inflammation), and no worsening in the potential remaining domain (defined as worsening  $\geq 20\%$  and  $\geq 1$  unit for ASAS20 or defined as worsening of  $> 0$  units for ASAS40).

<sup>b</sup> primary endpoint

<sup>c</sup> multiplicity-controlled  $p \leq 0.001$  upadacitinib vs placebo comparison

<sup>d</sup> multiplicity-controlled  $p \leq 0.01$  upadacitinib vs placebo comparison

<sup>e</sup> comparison not multiplicity-controlled

<sup>f</sup> post-hoc analysis for SELECT-AXIS 1, not multiplicity-controlled

For binary endpoints, week 14 results are based on non-responder imputation (SELECT-AXIS 1) and on non-responder imputation in conjunction with multiple imputation (SELECT-AXIS 2 [AS]). For continuous endpoints, week 14 results are based on the least squares mean change from baseline using mixed models for repeated measures analysis.

In SELECT-AXIS 1, efficacy was maintained through 2 years as assessed by the endpoints presented in Table 9.

#### *Physical function response and health-related outcomes*

In both studies, patients treated with upadacitinib 15 mg showed significant improvement in physical function compared to placebo as assessed by the Bath Ankylosing Spondylitis Functional Index (BASFI) change from baseline at week 14. In SELECT-AXIS 1, improvement in BASFI was maintained through 2 years.

In SELECT-AXIS 2 (AS), patients treated with upadacitinib 15 mg showed significant improvements in total back pain and nocturnal back pain compared to placebo at week 14.

In SELECT-AXIS 2 (AS), patients treated with upadacitinib 15 mg showed significant improvements in health-related quality of life and overall health as measured by ASQoL and ASAS Health Index, respectively, compared to placebo at week 14.

#### *Enthesitis*

In SELECT-AXIS 2 (AS), patients with pre-existing enthesitis (n=310) treated with upadacitinib 15 mg showed significant improvement in enthesitis compared to placebo as measured by change from baseline in Maastricht Ankylosing Spondylitis Enthesitis Score (MASSES) at week 14.

#### *Spinal mobility*

In SELECT-AXIS 2 (AS), patients treated with upadacitinib 15 mg showed significant improvement in spinal mobility compared to placebo as measured by change from baseline in Bath Ankylosing Spondylitis Metrology Index (BASMI) at week 14.

#### *Objective measure of inflammation*

Signs of inflammation were assessed by MRI and expressed as change from baseline in the SPARCC score for spine. In both studies, at week 14, significant improvement of inflammatory signs in the

spine was observed in patients treated with upadacitinib 15 mg compared to placebo. In SELECT-AXIS 1, improvement in inflammation as assessed by MRI was maintained through 2 years.

### *Atopic dermatitis*

The efficacy and safety of upadacitinib 15 mg and 30 mg once daily was assessed in three Phase 3 randomised, double-blind, multicentre studies (MEASURE UP 1, MEASURE UP 2 and AD UP) in a total of 2584 patients (12 years of age and older). Upadacitinib was evaluated in 344 adolescent and 2240 adult patients with moderate to severe atopic dermatitis (AD) not adequately controlled by topical medication(s). At baseline, patients had to have all the following: an Investigator's Global Assessment (vIGA-AD) score  $\geq 3$  in the overall assessment of AD (erythema, induration/papulation, and oozing/crusting) on an increasing severity scale of 0 to 4, an Eczema Area and Severity Index (EASI) score  $\geq 16$  (composite score assessing extent and severity of erythema, oedema/papulation, scratches and lichenification across 4 different body sites), a minimum body surface area (BSA) involvement of  $\geq 10\%$ , and weekly average Worst Pruritus Numerical Rating Scale (NRS)  $\geq 4$ .

In all three studies, patients received upadacitinib once daily doses of 15 mg, 30 mg, or matching placebo for 16 weeks. In the AD UP study, patients also received concomitant topical corticosteroids (TCS). Following completion of the double blinded period, patients originally randomised to upadacitinib were to continue receiving the same dose until week 260. Patients in the placebo group were re-randomised in a 1:1 ratio to receive upadacitinib 15 mg or 30 mg until week 260.

### *Baseline characteristics*

In the monotherapy studies (MEASURE UP 1 and 2), 50.0% of patients had a baseline vIGA-AD score of 3 (moderate) and 50.0% of patients had a baseline vIGA-AD of 4 (severe). The mean baseline EASI score was 29.3 and the mean baseline weekly average Worst Pruritus NRS was 7.3. In the concomitant TCS study (AD UP), 47.1% of patients had a baseline vIGA-AD score of 3 (moderate) and 52.9% of patients had a baseline vIGA-AD of 4 (severe). The mean baseline EASI score was 29.7 and the mean baseline weekly average Worst Pruritus NRS was 7.2.

### *Clinical response*

#### *Monotherapy (MEASURE UP 1 AND MEASURE UP 2) and Concomitant TCS (AD UP) studies*


A significantly greater proportion of patients treated with upadacitinib 15 mg or 30 mg achieved vIGA-AD 0 or 1, EASI 75, or a  $\geq 4$ -point improvement on the Worst Pruritus NRS compared to placebo at week 16. Rapid improvements in skin clearance and itch were also achieved (see Table [4011](#)).

Figure 1 shows the proportion of patients achieving an EASI 75 response and mean percent change from baseline in Worst Pruritus NRS, respectively up to week 16 for MEASURE UP 1 and 2.

**Table 4011: Efficacy results of upadacitinib**

Study	MEASURE UP 1			MEASURE UP 2			AD UP		
	PBO	UPA 15 mg	UPA 30 mg	PBO	UPA 15 mg	UPA 30 mg	PBO + TCS	UPA 15 mg + TCS	UPA 30 mg + TCS
Number of subjects randomised	281	281	285	278	276	282	304	300	297
<b>Week 16 endpoints, % responders (95% CI)</b>									
vIGA-AD 0/1 <sup>a,b</sup> (co-primary)	8 (5,12)	48 <sup>d</sup> (42,54)	62 <sup>d</sup> (56,68)	5 (2,7)	39 <sup>d</sup> (33,45)	52 <sup>d</sup> (46,58)	11 (7,14)	40 <sup>d</sup> (34,45)	59 <sup>d</sup> (53,64)
EASI 75 <sup>a</sup> (co-primary)	16 (12,21)	70 <sup>d</sup> (64,75)	80 <sup>d</sup> (75,84)	13 (9,17)	60 <sup>d</sup> (54,66)	73 <sup>d</sup> (68,78)	26 (21,31)	65 <sup>d</sup> (59,70)	77 <sup>d</sup> (72,82)
EASI 90 <sup>a</sup>	8 (5,11)	53 <sup>d</sup> (47,59)	66 <sup>d</sup> (60,71)	5 (3,8)	42 <sup>d</sup> (37,48)	58 <sup>d</sup> (53,64)	13 (9,17)	43 <sup>d</sup> (37,48)	63 <sup>d</sup> (58,69)
EASI 100 <sup>a</sup>	2 (0,3)	17 <sup>d</sup> (12,21)	27 <sup>d</sup> (22,32)	1 (0,2)	14 <sup>d</sup> (10,18)	19 <sup>d</sup> (14,23)	1 (0,3)	12 <sup>e</sup> (8,16)	23 <sup>d</sup> (18,27)
Worst Pruritus NRS <sup>c</sup> (≥ 4-point improvement)	12 (8,16)	52 <sup>d</sup> (46,58)	60 <sup>d</sup> (54,66)	9 (6,13)	42 <sup>d</sup> (36,48)	60 <sup>d</sup> (54,65)	15 (11,19)	52 <sup>d</sup> (46,58)	64 <sup>d</sup> (58,69)
<b>Early onset endpoints, % responders (95% CI)</b>									
EASI 75 <sup>a</sup> (Week 2)	4 (1,6)	38 <sup>d</sup> (32,44)	47 <sup>d</sup> (42,53)	4 (1,6)	33 <sup>d</sup> (27,39)	44 <sup>d</sup> (38,50)	7 (4,10)	31 <sup>d</sup> (26,36)	44 <sup>d</sup> (38,50)
Worst Pruritus NRS (≥ 4-point improvement at week 1) <sup>e,f</sup>	0 (0,1)	15 <sup>d</sup> (11,19)	20 <sup>d</sup> (15,24)	1 (0,2)	7 <sup>d</sup> (4,11)	16 <sup>d</sup> (11,20)	3 (1,5)	12 <sup>d</sup> (8,16)	19 <sup>d</sup> (15,24)
<p>Abbreviations: UPA= upadacitinib (RINVOQ); PBO = placebo</p> <p>Subjects with rescue medication or with missing data were counted as non-responders. The number and percentage of subjects who were imputed as non-responders for EASI 75 and vIGA-AD 0/1 at Week 16 due to the use of rescue therapy in the placebo, upadacitinib 15 mg, and upadacitinib 30 mg groups, respectively, were 132 (47.0%), 31 (11.0%), 16 (5.6%) in MEASURE UP 1, 119 (42.8%), 24 (8.7%), 16 (5.7%) in MEASURE UP 2, and 78 (25.7%), 15 (5.0%), 14 (4.7%) in AD UP.</p> <p><sup>a</sup> Based on number of subjects randomised</p> <p><sup>b</sup> Responder was defined as a patient with vIGA-AD 0 or 1 (“clear” or “almost clear”) with a reduction of ≥ 2 points on a 0-4 ordinal scale</p> <p><sup>c</sup> Results shown in subset of patients eligible for assessment (patients with Worst Pruritus NRS ≥ 4 at baseline)</p> <p><sup>d</sup> Statistically significant vs. placebo with p &lt; 0.001</p> <p><sup>e</sup> p &lt; 0.001 vs placebo, without multiplicity control</p> <p><sup>f</sup> Statistically significant improvements vs placebo were seen as early as 1 day after initiating upadacitinib 30 mg and 2 days after initiating upadacitinib 15 mg in MEASURE UP 1 and 2</p>									

**Figure 1:** Proportion of patients achieving an EASI 75 response and mean percent change from baseline in Worst Pruritus NRS in MEASURE UP 1 and MEASURE UP 2



Treatment effects in subgroups (weight, age, gender, race, and prior systemic treatment with immunosuppressants) were consistent with the results in the overall study population.

Results at week 16 continued to be maintained through week 52 in patients treated with upadacitinib 15 mg or 30 mg.

*Quality of life/patient-reported outcomes*

**Table 12:** Patient-reported outcomes results of upadacitinib at week 16

Study	MEASURE UP 1			MEASURE UP 2		
	PBO	UPA 15 mg	UPA 30 mg	PBO	UPA 15 mg	UPA 30 mg
Treatment group	PBO	UPA 15 mg	UPA 30 mg	PBO	UPA 15 mg	UPA 30 mg
Number of subjects randomised	281	281	285	278	276	282
<b>% responders (95% CI)</b>						
ADerm-SS Skin Pain (≥ 4-point improvement) <sup>a</sup>	15 (10,20)	54 <sup>e</sup> (47,60)	63 <sup>e</sup> (57,69)	13 (9,18)	49 <sup>e</sup> (43,56)	65 <sup>e</sup> (59,71)
ADerm-IS Sleep (≥ 12-point improvement) <sup>a,b</sup>	13 (9,18)	55 <sup>e</sup> (48,62)	66 <sup>e</sup> (60,72)	12 (8,17)	50 <sup>e</sup> (44,57)	62 <sup>e</sup> (56,69)
DLQI 0/1 <sup>c</sup>	4 (2,7)	30 <sup>e</sup> (25,36)	41 <sup>e</sup> (35,47)	5 (2,7)	24 <sup>e</sup> (19,29)	38 <sup>e</sup> (32,44)

Study	MEASURE UP 1			MEASURE UP 2		
	HADS Anxiety <8 and HADS Depression < 8 <sup>d</sup>	14 (8,20)	46 <sup>e</sup> (37,54)	49 <sup>e</sup> (41,57)	11 (6,17)	46 <sup>e</sup> (38,54)

Abbreviations: UPA= upadacitinib (RINVOQ); PBO = placebo; DLQI = Dermatology Life Quality Index; HADS = Hospital Anxiety and Depression Scale  
Subjects with rescue medication or with missing data were counted as non-responders.  
The threshold values specified correspond to the minimal clinically important difference (MCID) and was used to determine response.  
<sup>a</sup> Results shown in subset of patients eligible for assessment (patients with assessment score > MCID at baseline).  
<sup>b</sup> ADerm-IS Sleep assesses difficulty falling asleep, sleep impact, and waking up at night due to AD.  
<sup>c</sup> Results shown in subset of patients eligible for assessment (patients with DLQI > 1 at baseline).  
<sup>d</sup> Results shown in subset of patients eligible for assessment (patients with HADS Anxiety ≥ 8 or HADS Depression ≥ 8 at baseline)  
<sup>e</sup> Statistically significant vs. placebo with p < 0.001

### Ulcerative colitis

The efficacy and safety of upadacitinib was evaluated in three multicentre, double-blind, placebo-controlled Phase 3 clinical studies: two replicate induction studies, UC-1 (U-ACHIEVE Induction) and UC-2 (U-ACCOMPLISH), and a maintenance study UC-3 (U-ACHIEVE Maintenance).

Disease activity was based on the adapted Mayo score (aMS, Mayo scoring system excluding Physician's Global Assessment), which ranged from 0 to 9 and has three subscores that were each scored 0 (normal) to 3 (most severe): stool frequency subscore (SFS), rectal bleeding subscore (RBS) and a centrally-reviewed endoscopy subscore (ES).

#### Induction studies (UC-1 and UC-2)

In UC-1 and UC-2, 988 patients (473 and 515 patients, respectively) were randomised to upadacitinib 45 mg once daily or placebo for 8 weeks with a 2:1 treatment allocation ratio and included in the efficacy analysis. All enrolled patients had moderately to severely active ulcerative colitis defined as an aMS of 5 to 9 with an ES of 2 or 3 and demonstrated prior treatment failure including inadequate response, loss of response, or intolerance to prior conventional and/or biologic treatment. Prior treatment failure to at least 1 biologic therapy (prior biologic failure) was seen in 52% (246/473) and 51% (262/515) of patients, respectively. Previous treatment failure to conventional therapy but not biologics (without prior biologic failure) was seen in 48% (227/473) and 49% (253/515) of patients, respectively.

At baseline in UC-1 and UC-2, 39% and 37% of patients received corticosteroids, 1.1% and 0.6% of patients received methotrexate and 68% and 69% of patients received aminosalicylates. Concomitant use of thiopurine was not allowed during the studies. Patient disease activity was moderate (aMS ≥5, ≤7) in 61% and 60% of patients and severe (aMS >7) in 39% and 40% of patients.

The primary endpoint was clinical remission per aMS at week 8. Table 13 shows the primary and key secondary endpoints including clinical response, mucosal healing, histologic-endoscopic mucosal healing and deep mucosal healing.



**Table 13 Proportion of patients meeting primary and key secondary efficacy endpoints at week 8 in the induction studies UC-1 and UC-2**

<u>Endpoint</u>	<u>UC-1 (U-ACHIEVE)</u>			<u>UC-2 (U-ACCOMPLISH)</u>		
	<u>PBO N=154</u>	<u>UPA 45 mg N=319</u>	<u>Treatment Difference (95% CI)</u>	<u>PBO N=174</u>	<u>UPA 45 mg N=341</u>	<u>Treatment Difference (95% CI)</u>
<u>Clinical remission<sup>a</sup></u>	<u>4.8%</u>	<u>26.1%</u>	<u>21.6%*</u> <u>(15.8, 27.4)</u>	<u>4.1%</u>	<u>33.5%</u>	<u>29.0%*</u> <u>(23.2, 34.7)</u>
<u>Prior biologic failure<sup>+</sup></u>	<u>0.4%</u>	<u>17.9%</u>	<u>17.5%</u>	<u>2.4%</u>	<u>29.6%</u>	<u>27.1%</u>
<u>Without prior biologic failure<sup>+</sup></u>	<u>9.2%</u>	<u>35.2%</u>	<u>26.0%</u>	<u>5.9%</u>	<u>37.5%</u>	<u>31.6%</u>
<u>Clinical response<sup>b</sup></u>	<u>27.3%</u>	<u>72.6%</u>	<u>46.3%*</u> <u>(38.4, 54.2)</u>	<u>25.4%</u>	<u>74.5%</u>	<u>49.4%*</u> <u>(41.7, 57.1)</u>
<u>Prior biologic failure<sup>+</sup></u>	<u>12.8%</u>	<u>64.4%</u>	<u>51.6%</u>	<u>19.3%</u>	<u>69.4%</u>	<u>50.1%</u>
<u>Without prior biologic failure<sup>+</sup></u>	<u>42.1%</u>	<u>81.8%</u>	<u>39.7%</u>	<u>31.8%</u>	<u>79.8%</u>	<u>48.0%</u>
<u>Mucosal healing<sup>c</sup></u>	<u>7.4%</u>	<u>36.3%</u>	<u>29.3%*</u> <u>(22.6, 35.9)</u>	<u>8.3%</u>	<u>44.0%</u>	<u>35.1%*</u> <u>(28.6, 41.6)</u>
<u>Prior biologic failure<sup>+</sup></u>	<u>1.7%</u>	<u>27.0%</u>	<u>25.3%</u>	<u>4.8%</u>	<u>37.1%</u>	<u>32.3%</u>
<u>Without prior biologic failure<sup>+</sup></u>	<u>13.2%</u>	<u>46.8%</u>	<u>33.6%</u>	<u>12.0%</u>	<u>51.2%</u>	<u>39.2%</u>
<u>Histologic-endoscopic mucosal healing<sup>d</sup></u>	<u>6.6%</u>	<u>30.1%</u>	<u>23.7%*</u> <u>(17.5, 30.0)</u>	<u>5.9%</u>	<u>36.7%</u>	<u>30.1%*</u> <u>(24.1, 36.2)</u>
<u>Prior biologic failure<sup>+</sup></u>	<u>1.4%</u>	<u>22.7%</u>	<u>21.3%</u>	<u>4.6%</u>	<u>30.7%</u>	<u>26.1%</u>
<u>Without prior biologic failure<sup>+</sup></u>	<u>11.8%</u>	<u>38.2%</u>	<u>26.4%</u>	<u>7.2%</u>	<u>42.9%</u>	<u>35.7%</u>
<u>Deep mucosal healing<sup>e</sup></u>	<u>1.3%</u>	<u>10.7%</u>	<u>9.7%*</u> <u>(5.7, 13.7)</u>	<u>1.7%</u>	<u>13.5%</u>	<u>11.3%*</u> <u>(7.2, 15.3)</u>
<u>Prior biologic failure<sup>+</sup></u>	<u>0</u>	<u>6.5%</u>	<u>6.5%</u>	<u>1.1%</u>	<u>9.2%</u>	<u>8.1%</u>
<u>Without prior biologic failure<sup>+</sup></u>	<u>2.6%</u>	<u>15.4%</u>	<u>12.8%</u>	<u>2.4%</u>	<u>17.9%</u>	<u>15.5%</u>



Abbreviations: PBO = placebo; UPA= upadacitinib; aMS = adapted Mayo Score, based on the Mayo Scoring system (excluding Physician's Global Assessment), which ranged from 0 to 9 and has three subscores that were each scored 0 (normal) to 3 (most severe): stool frequency subscore (SFS), rectal bleeding subscore (RBS) and a centrally-reviewed endoscopy subscore (ES).

+The number of “Prior biologic failure” patients in UC-1 and UC-2 are 78 and 89 in the placebo group, and 168 and 173 in the upadacitinib 45 mg group, respectively; the number of “Without prior biologic failure” patients in UC-1 and UC-2 are 76 and 85 in the placebo group, and 151 and 168 in the upadacitinib 45 mg group, respectively.

\*p <0.001, adjusted treatment difference (95% CI)

<sup>a</sup> Per aMS: SFS < 1 and not greater than baseline, RBS = 0, ES < 1 without friability

<sup>b</sup> Per aMS: decrease ≥ 2 points and ≥ 30% from baseline and a decrease in RBS ≥ 1 from baseline or an absolute RBS ≤ 1.

<sup>c</sup> ES < 1 without friability

<sup>d</sup> ES ≤ 1 without friability and Geboes score ≤ 3.1 (indicating neutrophil infiltration in < 5% of crypts, no crypt destruction, and no erosions, ulcerations, or granulation tissue.)

<sup>e</sup> ES = 0, Geboes score < 2 (indicating no neutrophil in crypts or lamina propria and no increase in eosinophil, no crypt destruction, and no erosions, ulcerations, or granulation tissue)

### Disease activity and symptoms

The partial adapted Mayo score (paMS) is composed of SFS and RBS. Symptomatic response per paMS is defined as a decrease of ≥1 point and ≥30% from baseline and a decrease in RBS ≥ 1 or an absolute RBS ≤1. Statistically significant improvement compared to placebo per paMS was seen as early as week 2 (UC-1: 60.1% vs 27.3% and UC-2: 63.3% vs 25.9%).

### Extended induction

A total of 125 patients in UC-1 and UC-2 who did not achieve clinical response after 8 weeks of treatment with upadacitinib 45 mg once daily entered an 8-week open-label extended induction period. After the treatment of an additional 8 weeks (16 weeks total) of upadacitinib 45 mg once daily, 48.3% of patients achieved clinical response per aMS. Among patients who responded to treatment of 16-week upadacitinib 45 mg once daily, 35.7% and 66.7% of patients maintained clinical response per aMS and 19.0% and 33.3% of patients achieved clinical remission per aMS at week 52 with maintenance treatment of upadacitinib 15 mg and 30 mg once daily, respectively.

### Maintenance study (UC-3)

The efficacy analysis for UC-3 was evaluated in 451 patients who achieved clinical response per aMS with 8-week upadacitinib 45 mg once daily induction treatment. Patients were randomised to receive upadacitinib 15 mg, 30 mg or placebo once daily for up to 52 weeks.

The primary endpoint was clinical remission per aMS at week 52. Table 14 shows the key secondary endpoints including maintenance of clinical remission, corticosteroid-free clinical remission, mucosal healing, histologic-endoscopic mucosal healing and deep mucosal healing.

**Table 14 Proportion of patients meeting primary and key secondary efficacy endpoints at week 52 in the maintenance study UC-3**

	<u>PBO</u> <u>N=149</u>	<u>UPA</u> <u>15 mg</u> <u>N=148</u>	<u>UPA</u> <u>30 mg</u> <u>N=154</u>	<u>Treatment</u> <u>Difference</u> <u>15 mg vs PBO</u> <u>(95% CI)</u>	<u>Treatment</u> <u>Difference</u> <u>30 mg vs PBO</u> <u>(95% CI)</u>
<u>Clinical remission<sup>a</sup></u>	<u>12.1%</u>	<u>42.3%</u>	<u>51.7%</u>	<u>30.7%*</u> <u>(21.7, 39.8)</u>	<u>39.0%*</u> <u>(29.7, 48.2)</u>
<u>Prior biologic failure<sup>+</sup></u>	<u>7.5%</u>	<u>40.5%</u>	<u>49.1%</u>	<u>33.0%</u>	<u>41.6%</u>
<u>Without prior biologic failure<sup>+</sup></u>	<u>17.6%</u>	<u>43.9%</u>	<u>54.0%</u>	<u>26.3%</u>	<u>36.3%</u>

<b><u>Maintenance of clinical remission<sup>b</sup></u></b>	<u>N = 54</u> <u>22.2%</u>	<u>N = 47</u> <u>59.2%</u>	<u>N = 58</u> <u>69.7%</u>	<u>37.4%*</u> <u>(20.3, 54.6)</u>	<u>47.0%*</u> <u>(30.7, 63.3)</u>
<u>Prior biologic failure</u>	<u>N = 22</u> <u>13.6%</u>	<u>N = 17</u> <u>76.5%</u>	<u>N = 20</u> <u>73.0%</u>	<u>62.8%</u>	<u>59.4%</u>
<u>Without prior biologic failure</u>	<u>N = 32</u> <u>28.1%</u>	<u>N = 30</u> <u>49.4%</u>	<u>N = 38</u> <u>68.0%</u>	<u>21.3%</u>	<u>39.9%</u>
<b><u>Corticosteroid-free clinical remission<sup>c</sup></u></b>	<u>N = 54</u> <u>22.2%</u>	<u>N = 47</u> <u>57.1%</u>	<u>N = 58</u> <u>68.0%</u>	<u>35.4%*</u> <u>(18.2, 52.7)</u>	<u>45.1%*</u> <u>(28.7, 61.6)</u>
<u>Prior biologic failure</u>	<u>N = 22</u> <u>13.6%</u>	<u>N = 17</u> <u>70.6%</u>	<u>N = 20</u> <u>73.0%</u>	<u>57.0%</u>	<u>59.4%</u>
<u>Without prior biologic failure</u>	<u>N = 32</u> <u>28.1%</u>	<u>N = 30</u> <u>49.4%</u>	<u>N = 38</u> <u>65.4%</u>	<u>21.3%</u>	<u>37.2%</u>
<b><u>Mucosal healing<sup>d</sup></u></b>	<u>14.5%</u>	<u>48.7%</u>	<u>61.6%</u>	<u>34.4%*</u> <u>(25.1, 43.7)</u>	<u>46.3%*</u> <u>(36.7, 55.8)</u>
<u>Prior biologic failure<sup>+</sup></u>	<u>7.8%</u>	<u>43.3%</u>	<u>56.1%</u>	<u>35.5%</u>	<u>48.3%</u>
<u>Without prior biologic failure<sup>+</sup></u>	<u>22.5%</u>	<u>53.6%</u>	<u>66.6%</u>	<u>31.1%</u>	<u>44.1%</u>
<b><u>Histologic-endoscopic mucosal healing<sup>e</sup></u></b>	<u>11.9%</u>	<u>35.0%</u>	<u>49.8%</u>	<u>23.8%*</u> <u>(14.8, 32.8)</u>	<u>37.3%*</u> <u>(27.8, 46.8)</u>
<u>Prior biologic failure<sup>+</sup></u>	<u>5.2%</u>	<u>32.9%</u>	<u>47.6%</u>	<u>27.7%</u>	<u>42.4%</u>
<u>Without prior biologic failure<sup>+</sup></u>	<u>20.0%</u>	<u>36.9%</u>	<u>51.8%</u>	<u>16.9%</u>	<u>31.8%</u>
<b><u>Deep mucosal healing<sup>f</sup></u></b>	<u>4.7%</u>	<u>17.6%</u>	<u>19.0%</u>	<u>13.0%*</u> <u>(6.0, 20.0)</u>	<u>13.6%*</u> <u>(6.6, 20.6)</u>
<u>Prior biologic failure<sup>+</sup></u>	<u>2.5%</u>	<u>17.2%</u>	<u>16.1%</u>	<u>14.7%</u>	<u>13.6%</u>
<u>Without prior biologic failure<sup>+</sup></u>	<u>7.5%</u>	<u>18.0%</u>	<u>21.6%</u>	<u>10.6%</u>	<u>14.2%</u>

Abbreviations: PBO = placebo; UPA= upadacitinib; aMS = adapted Mayo Score, based on the Mayo Scoring system (excluding Physician's Global Assessment), which ranged from 0 to 9 and has three subscores that were each scored 0 (normal) to 3 (most severe): stool frequency subscore (SFS), rectal bleeding subscore (RBS) and a centrally-reviewed endoscopy subscore (ES).

<sup>+</sup>The number of “Prior biologic failure” patients are 81, 71, and 73 in the placebo, upadacitinib 15 mg, and 30 mg group, respectively. The number of “Without prior biologic failure” patients are 68, 77, and 81 in the placebo, upadacitinib 15 mg, and 30 mg group, respectively.

\* p <0.001, adjusted treatment difference (95% CI)

<sup>a</sup> Per aMS: SFS ≤ 1 and not greater than baseline, RBS = 0, ES ≤ 1 without friability

<sup>b</sup> Clinical remission per aMS at Week 52 among patients who achieved clinical remission at the end of induction treatment.

<sup>c</sup> Clinical remission per aMS at Week 52 and corticosteroid-free for >90 days immediately preceding Week 52 among patients who achieved clinical remission at the end of the induction treatment.

<sup>d</sup> ES ≤1 without friability


<sup>e</sup> ES ≤1 without friability and Geboes score ≤ 3.1 (indicating neutrophil infiltration in <5% of crypts, no crypt destruction and no erosions, ulcerations or granulation tissue).

<sup>f</sup> ES = 0, Geboes score < 2 (indicating no neutrophil in crypts or lamina propria and no increase in eosinophil, no crypt destruction, and no erosions, ulcerations or granulation tissue).

### Disease symptoms

Symptomatic remission per paMS, defined as SFS ≤ 1 and RBS = 0, was achieved over time through week 52 in more patients treated with both upadacitinib 15 mg and 30 mg once daily compared with placebo (Figure 2).

### Figure 2 Proportion of patients with symptomatic remission per partial adapted Mayo score over time in maintenance study UC-3



### Endoscopic assessment

Endoscopic remission (normalisation of the endoscopic appearance of the mucosa) was defined as ES of 0. At week 8, a significantly greater proportion of patients treated with upadacitinib 45 mg once daily compared to placebo achieved endoscopic remission (UC-1: 13.7% vs 1.3%, UC-2: 18.2% vs 1.7%). In UC-3, a significantly greater proportion of patients treated with upadacitinib 15 mg and 30 mg once daily compared to placebo achieved endoscopic remission at week 52 (24.2% and 25.9% vs 5.6%). Maintenance of mucosal healing at week 52 (ES ≤1 without friability) was seen in a significantly greater proportion of patients treated with upadacitinib 15 mg and 30 mg once daily compared to placebo (61.6% and 69.5% vs 19.2%) among patients who achieved mucosal healing at the end of induction.

### Quality of life

Patients treated with upadacitinib demonstrated significantly greater and clinically meaningful improvement in health-related quality of life measured by the Inflammatory Bowel Disease Questionnaire (IBDQ) total score compared to placebo. Improvements were seen in all 4 domain scores: systemic symptoms (including fatigue), social function, emotional function and bowel symptoms (including abdominal pain and bowel urgency). Changes in IBDQ total score at week 8 from baseline with upadacitinib 45 mg once daily compared to placebo were 55.3 and 21.7 in UC-1 and 52.2 and 21.1 in UC-2, respectively. Changes in IBDQ total score at week 52 from baseline were 49.2, 58.9 and 17.9 in patients treated with upadacitinib 15 mg, 30 mg once daily and placebo, respectively.

### Paediatric population

A total of 344 adolescents aged 12 to 17 years with moderate to severe atopic dermatitis were randomised across the three Phase 3 studies to receive either 15 mg (N=114) or 30 mg (N=114) upadacitinib or matching placebo (N=116), in monotherapy or combination with topical corticosteroids. Efficacy was consistent between the adolescents and adults. The safety profile in adolescents was generally similar to that in adults, with dose dependent increases in the rate of some adverse events, including neutropaenia and herpes zoster. At both doses, the rate of neutropaenia was slightly increased in adolescents compared to adults. The rate of herpes zoster in adolescents at the 30 mg dose was comparable to that in adults. The safety and efficacy of the 30 mg dose in adolescents are still being investigated.

**Table 12:15 Efficacy results of upadacitinib for adolescents at week 16**

Study	MEASURE UP 1		MEASURE UP 2		AD UP	
	PBO	UPA 15 mg	PBO	UPA 15 mg	PBO + TCS	UPA 15 mg + TCS
Number of adolescent subjects randomised	40	42	36	33	40	39
<b>% responders (95% CI)</b>						
vIGA-AD 0/1 <sup>a,b</sup>	8 (0,16)	38 (23,53)	3 (0,8)	42 (26,59)	8 (0,16)	31 (16,45)
EASI 75 <sup>a</sup>	8 (0,17)	71 (58,85)	14 (3,25)	67 (51,83)	30 (16,44)	56 (41,72)
Worst Pruritus NRS <sup>c</sup> (≥ 4-point improvement)	15 (4,27)	45 (30,60)	3 (0,8)	33 (16,50)	13 (2,24)	42 (26,58)
Abbreviations: UPA= upadacitinib (RINVOQ); PBO = placebo Subjects with rescue medication or with missing data were counted as non-responders. <sup>a</sup> Based on number of subjects randomised <sup>b</sup> Responder was defined as a patient with vIGA-AD 0 or 1 (“clear” or “almost clear”) with a reduction of ≥ 2 points on a 0-4 ordinal scale. <sup>c</sup> Results shown in subset of patients eligible for assessment (patients with Worst Pruritus NRS ≥ 4 at baseline).						

## 5.2 Pharmacokinetic properties

Upadacitinib plasma exposures are proportional to dose over the therapeutic dose range. Steady-state plasma concentrations are achieved within 4 days with minimal accumulation after multiple once-daily administrations.

### *Absorption*

Following oral administration of upadacitinib prolonged-release formulation, upadacitinib is absorbed with a median  $T_{max}$  of 2 to 4 hours. Coadministration of upadacitinib with a high-fat meal had no clinically relevant effect on upadacitinib exposures (increased AUC by 29% and  $C_{max}$  by 39% **to 60%**). In clinical trials, upadacitinib was administered without regard to meals (see section 4.2). *In vitro*, upadacitinib is a substrate for the efflux transporters P-gp and BCRP.

### *Distribution*

Upadacitinib is 52% bound to plasma proteins. Upadacitinib partitions similarly between plasma and blood cellular components, as indicated by the blood to plasma ratio of 1.0.

### *Metabolism*

Upadacitinib metabolism is mediated by CYP3A4 with a potential minor contribution from CYP2D6. The pharmacologic activity of upadacitinib is attributed to the parent molecule. In a human radiolabeled study, unchanged upadacitinib accounted for 79% of the total radioactivity in plasma while the main metabolite (product of monooxidation followed by glucuronidation) accounted for 13% of the total plasma radioactivity. No active metabolites have been identified for upadacitinib.

### *Elimination*

Following single dose administration of [<sup>14</sup>C]-upadacitinib immediate-release solution, upadacitinib was eliminated predominantly as the unchanged parent substance in urine (24%) and faeces (38%). Approximately 34% of upadacitinib dose was excreted as metabolites. Upadacitinib mean terminal elimination half-life ranged from 9 to 14 hours.

### Special populations

#### Renal impairment

Upadacitinib AUC was 18%, 33%, and 44% higher in subjects with mild (estimated glomerular filtration rate 60 to 89 mL/min/1.73 m<sup>2</sup>), moderate (estimated glomerular filtration rate 30 to 59 mL/min/1.73 m<sup>2</sup>), and severe (estimated glomerular filtration rate 15 to 29 mL/min/1.73 m<sup>2</sup>) renal impairment, respectively, compared to subjects with normal renal function. Upadacitinib C<sub>max</sub> was similar in subjects with normal and impaired renal function. Mild or moderate renal impairment has no clinically relevant effect on upadacitinib exposure ~~following the 15 mg or 30 mg once daily dosing regimens. The recommended dose is 15 mg once daily for patients with severe renal impairment.~~ (see [section 4.2](#)).

#### Hepatic impairment

Mild (Child-Pugh A) and moderate (Child-Pugh B) hepatic impairment has no clinically relevant effect on upadacitinib exposure. Upadacitinib AUC was 28% and 24% higher in subjects with mild and moderate hepatic impairment, respectively, compared to subjects with normal liver function. Upadacitinib C<sub>max</sub> was unchanged in subjects with mild hepatic impairment and 43% higher in subjects with moderate hepatic impairment compared to subjects with normal liver function. Upadacitinib was not studied in patients with severe (Child-Pugh C) hepatic impairment.

#### Paediatric population

The pharmacokinetics of upadacitinib have not yet been evaluated in paediatric patients with rheumatoid arthritis, psoriatic arthritis, ~~and~~ ankylosing spondylitis [and ulcerative colitis](#) (see section 4.2).

Upadacitinib pharmacokinetics and steady-state concentrations are similar for adults and adolescents 12 to 17 years of age with atopic dermatitis. The posology in adolescent patients 30 kg to < 40 kg was determined using population pharmacokinetic modelling and simulation.

The pharmacokinetics of upadacitinib in paediatric patients (< 12 years of age) with atopic dermatitis have not been established.

#### Intrinsic factors

Age, sex, body weight, race, and ethnicity did not have a clinically meaningful effect on upadacitinib exposure. Upadacitinib pharmacokinetics are consistent between rheumatoid arthritis, psoriatic arthritis, ankylosing spondylitis, ~~and~~ atopic dermatitis [and ulcerative colitis](#) patients.

### **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology.

Upadacitinib, at exposures (based on AUC) approximately 4 and 10 times the clinical dose of 15 mg, ~~and 2 and 5 times the clinical dose of 30 mg,~~ and 1.6 and 4 times the clinical dose of 45 mg in male and female Sprague-Dawley rats, respectively, was not carcinogenic in a 2-year carcinogenicity study in Sprague-Dawley rats. Upadacitinib was not carcinogenic in a 26-week carcinogenicity study in CByB6F1-Tg(HRAS)2Jic transgenic mice.

Upadacitinib was not mutagenic or genotoxic based on the results of *in vitro* and *in vivo* tests for gene mutations and chromosomal aberrations.

Upadacitinib had no effect on fertility in male or female rats at exposures up to approximately ~~24~~ 16 and ~~43~~ 31 times the maximum recommended human dose (MRHD) of ~~30~~ 45 mg in males and females, respectively, on an AUC basis in a fertility and early embryonic development study.

Dose-related increases in foetal resorptions associated with post-implantation losses in this fertility study in rats were attributed to the developmental/teratogenic effects of upadacitinib. No adverse effects were observed at exposures below clinical exposure (based on AUC). Post-implantation losses were observed at exposures ~~11~~ 8 times the clinical exposure at the MRHD of ~~30~~ 45 mg (based on AUC).

In animal embryo-foetal development studies, upadacitinib was teratogenic in both rats and rabbits. Upadacitinib resulted in increases in skeletal malformations in rats at 1.6, ~~and 0.8,~~ and 0.6 times the clinical exposure (AUC-based) at the 15, ~~and 30,~~ and 45 mg (MRHD) doses, respectively. In rabbits an increased incidence of cardiovascular malformations was observed at 15, ~~and 7.6~~ and 5.6 times the clinical exposure at the 15, ~~and 30,~~ and 45 mg doses (AUC-based), respectively. ~~No developmental toxicity was observed at approximately 0.15 times (rat) and at similar exposure in rabbits as the exposures at the MRHD of 30 mg. In a pre- and post-natal development study in pregnant female rats, oral upadacitinib administration at exposures approximately 1.4 times the MRHD of 30 mg resulted in no maternal effects, no effects on parturition, lactation or maternal behaviour and no effects on the offspring.~~

Following administration of upadacitinib to lactating rats, the concentrations of upadacitinib in milk over time generally paralleled those in plasma, with approximately 30-fold higher exposure in milk relative to maternal plasma. Approximately 97% of upadacitinib-related material in milk was the parent molecule, upadacitinib.

~~Administration of upadacitinib to juvenile Sprague-Dawley rats (from postnatal day 15 to 63) resulted in exposures and pharmacologic effects on the lymphoid system similar to those observed in adult rats. No adverse findings were observed in juvenile rats at exposures (AUC) approximately 9.4 and 4.8 times the exposures at the clinical doses of 15 mg and 30 mg, respectively (based on exposures in adult RA patients).~~

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

#### Tablet contents:

Microcrystalline cellulose  
Mannitol  
Hypromellose  
Tartaric acid (powdered)  
Magnesium stearate  
Silica, colloidal anhydrous / Colloidal Silicon Dioxide

#### Film coating:

Polyvinyl alcohol  
Macrogol /Polyethylene Glycol  
Talc  
Titanium dioxide (E171)  
Black Iron oxide (E172) / Ferrosferric Oxide (15 mg strength only)  
Iron oxide red (E172)

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

The expiry date of the product is indicated on the packaging materials.

## **6.4 Special precautions for storage**

Store up to 30°C.

Store in the original blister or bottle in order to protect from moisture. Keep the bottle tightly closed.

## **6.5 Nature and contents of container**

### RINVOQ 15 mg prolonged-release tablets

Polyvinylchloride/polyethylene/polychlorotrifluoroethylene - aluminium calendar blisters in packs containing 28 or 98 prolonged-release tablets, or multipacks containing 84 (3 packs of 28) prolonged-release tablets.

HDPE bottles with desiccant and polypropylene cap in carton containing 30 prolonged-release tablets. Pack size: 1 bottle (30 prolonged-release tablets) or 3 bottles (90 prolonged-release tablets).

Not all pack sizes may be marketed.

### RINVOQ 30 mg prolonged-release tablets

Polyvinylchloride/polyethylene/polychlorotrifluoroethylene - aluminium calendar blisters in packs containing 28 or 98 prolonged-release tablets.

HDPE bottles with desiccant and polypropylene cap in carton containing 30 prolonged-release tablets. Pack size: 1 bottle (30 prolonged-release tablets) or 3 bottles (90 prolonged-release tablets).

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MANUFACTURER**

AbbVie Deutschland GmbH & Co. KG, Knollstrasse, 67061 Ludwigshafen, Germany

## **8. LICENSE HOLDER**

AbbVie biopharmaceuticals LTD., 4 Hacharash St., Hod Hasharon, Israel.

**9. REGISTRATION NUMBER**

164-29-36194

168-69-37036

Revised on ~~July~~ August 2022 Based on MoH guidelines



## עלון לצרכן לפי תקנות הרוקחים (תכשירים) התשמ"ו – 1986

התרופה משווקת על-פי מרשם רופא בלבד

רינבוק<sup>TM</sup> 15 מ"ג

רינבוק<sup>TM</sup> 30 מ"ג

טבליות לשחרור ממושך

### המרכיב הפעיל וכמותו:

כל טבליה לשחרור ממושך מכילה אופדסיטיניב (כהמיהדרט) 15 מ"ג

Upadacitinib (as hemihydrate) 15mg

כל טבליה לשחרור ממושך מכילה אופדסיטיניב (כהמיהדרט) 30 מ"ג

Upadacitinib (as hemihydrate) 30mg

**קרא בעיון את העלון עד סופו בטרם תשתמש בתרופה.** עלון זה מכיל מידע תמציתי על התרופה. אם יש לך שאלות נוספות, פנה אל הרופא או אל הרוקח.  
תרופה זו נרשמה לטיפול עבורך. אל תעביר אותה לאחרים. היא עלולה להזיק להם אפילו אם נראה לך כי מצבם הרפואי דומה.

בנוסף לעלון, לתכשיר רינבוק קיים 'כרטיס מידע בטיחותי למטופל'. כרטיס זה מכיל מידע בטיחותי חשוב, שעליך לדעת, לפני התחלת הטיפול ובמהלך הטיפול ברינבוק ולפעול על פיו. יש לעיין ב'כרטיס מידע בטיחותי למטופל' ובעלון לצרכן בטרם תחילת השימוש בתכשיר. יש לשמור את הכרטיס לעיון נוסף במידת הצורך.

### 1. למה מיועדת התרופה?

רינבוק מיועד לטיפול ב:

- דלקת מפרקים שגרונית (Rheumatoid Arthritis) בינונית עד חמורה במבוגרים, אשר חוו תגובה לא מספקת בטיפול קודם בתרופה אנטי ראומטית אחת או יותר מקבוצת DMARDs או שהטיפול היה בלתי נסבל עבורם. ניתן להשתמש ברינבוק כטיפול יחיד או בשילוב עם מתוטרקסאט (methotrexate).
- דלקת מפרקים ספחתית (Psoriatic Arthritis) במבוגרים, אשר חוו תגובה לא מספקת בטיפול קודם בתרופה אנטי ראומטית אחת או יותר מקבוצת DMARDs או שהטיפול היה בלתי נסבל עבורם. ניתן להשתמש ברינבוק כטיפול יחיד או בשילוב עם מתוטרקסאט (methotrexate).
- דלקת חוליות מקשחת (Ankylosing Spondylitis) במבוגרים, אשר חוו תגובה לא מספקת לטיפול מקובל.
- דלקת עור אטופית (Atopic Dermatitis) בינונית עד חמורה במבוגרים ובמתבגרים בגילאי 12 ומעלה אשר מועמדים לטיפול סיסטמי.
- קוליטיס כיבית (Ulcerative Colitis) בינונית עד חמורה במבוגרים, אשר חוו תגובה לא מספקת, אובדן תגובה או אי סבילות לטיפול קונבנציונלי או ביולוגי קודם.

**קבוצה תרפויטית:** מדכאי מערכת חיסון סלקטיביים, מעכבי ג'אנוס קינאז.

רינבוק פועל על ידי הפחתת הפעילות של אנזימים בגוף הנקרא "ג'אנוס קינאז" אשר מוריד דלקת.

### דלקת מפרקים שגרונית

רינבוק משמש לטיפול בדלקת מפרקים שגרונית במבוגרים. דלקת מפרקים שגרונית זוהי מחלה הגורמת למפרקים

מודלקים. אם אתה סובל מדלקת מפרקים שגרונת פעילה בינונית עד חמורה, ייתכן ותחילה תקבל תרופות אחרות, אשר אחת מהן תהיה ככל הנראה מתוטרקסאט. אם תרופות אלו לא השפיעו באופן מספק, תקבל רינבוק עם או ללא מתוטרקסאט לטיפול בדלקת המפרקים השגרונת שלך.

רינבוק יכול לעזור בהפחתת כאב, נוקשות ונפיחות במפרקים שלך, להפחית עייפות, ויכול להאט את הנזק לעצם ולסחוס במפרקים שלך. השפעות אלו יכולות להקל על הפעילות היום-יומית הרגילה שלך ובכך לשפר את איכות חייך.

#### **דלקת מפרקים ספחתית**

רינבוק משמש לטיפול בדלקת מפרקים ספחתית במבוגרים. דלקת מפרקים ספחתית הינה מחלה הגורמת למפרקים מודלקים ופסוריאזיס. אם אתה סובל מדלקת מפרקים ספחתית פעילה, ייתכן ותחילה תקבל תרופות אחרות. אם תרופות אלו לא השפיעו באופן מספק, תקבל רינבוק עם או ללא מתוטרקסאט לטיפול בדלקת המפרקים הספחתית שלך.

רינבוק יכול לעזור בהפחתת כאב, נוקשות ונפיחות במפרקים שלך ומסביבם, כאב ונוקשות בעמוד השדרה, נגעים ספחתיים בעור ועייפות ויכול להאט את הנזק לעצם ולסחוס במפרקים שלך. השפעות אלו יכולות להקל על הפעילות היום-יומית הרגילה שלך ובכך לשפר את איכות חייך.

#### **דלקת חוליות מקשחת**

רינבוק משמש לטיפול בדלקת חוליות מקשחת במבוגרים. דלקת חוליות מקשחת היא מחלה הגורמת בעיקר לדלקת בעמוד השדרה. אם אתה סובל מדלקת חוליות מקשחת פעילה, ייתכן ותחילה תקבל תרופות אחרות. אם תרופות אלו לא השפיעו באופן מספק, תקבל רינבוק לטיפול בדלקת החוליות המקשחת שלך.

רינבוק יכול לעזור בהפחתת כאבי גב, נוקשות ודלקת בעמוד השדרה. השפעות אלו יכולות להקל על הפעילות היום-יומית שלך ובכך לשפר את איכות חייך.

#### **דלקת עור אטופית**

רינבוק משמש לטיפול בדלקת עור אטופית בינונית עד חמורה במבוגרים ובמתבגרים בגילאי 12 ומעלה. רינבוק ניתן לשימוש עם תרופות למריחה על העור שניתנות לדלקת עור אטופית או כטיפול יחיד.

נטילת רינבוק יכולה לשפר את מצב העור, להפחית גרד והתלקחויות. רינבוק יכול לסייע בשיפור תסמינים של כאב, חרדה ודיכאון אשר יכולים להופיע בחולים עם דלקת עור אטופית. רינבוק יכול גם לסייע בשיפור הפרעות שינה ואיכות החיים הכוללת.

#### **קוליטיס כיבית**

קוליטיס כיבית זו מחלה דלקתית של המעי הגס. רינבוק משמשת לטיפול במבוגרים הסובלים מקוליטיס כיבית אשר הגיבו בצורה לא מספקת או שהטיפול לא היה נסבל עבורם.

רינבוק יכול לעזור בהפחתת סימנים ותסמינים של המחלה הכוללים דימום צואתי, כאב בטני דחיפות ותכיפות הליכה לשירותים. השפעות אלו יכולות לאפשר קיום שגרת חיים תקינה והקלה בתשיות.

## 2. לפני השימוש בתרופה

### אין להשתמש בתרופה אם:

- אתה רגיש (אלרגי) לאופדסיטיניב או לכל אחד מהמרכיבים הנוספים אשר מכילה התרופה (מפורטים בסעיף 6)
- יש לך זיהום חריף (כגון דלקת ריאות או זיהום עור חיידקי)
- יש לך שחפת פעילה (TB)
- יש לך בעיות כבד חמורות
- הינך בהריון (ראי סעיף "הריון, הנקה ופוריות")

### אזהרות מיוחדות הנוגעות לשימוש בתרופה

#### לפני הטיפול ובמהלך הטיפול ברינבוק ספר לרופא אם:

- יש לך זיהום (חום, הזעה, או צמרמורות, קוצר נשימה, עור חם, אדום או כואב או פצעים על גופך, הרגשת עייפות, שיעול, תחושת צריבה בעת מתן שתן או מתן שתן לעיתים תכופות יותר מהרגיל, כאב ראש חמור עם צוואר נוקשה), או אם אי פעם היה לך זיהום חוזר – רינבוק יכול להפחית את יכולת הגוף שלך להילחם בזיהומים ולכן עשוי להחמיר זיהום קיים, או להגביר את הסיכוי שלך לפתח זיהום חדש
- סבלת משחפת או שהיית במגע קרוב עם מישהו חולה שחפת. הרופא שלך יבצע בדיקה לשחפת לפני התחלת הטיפול ברינבוק וייתכן שיחזור על הבדיקה בזמן הטיפול
- סבלת מזיהום הרפס זוסטר (שלבקת חוגרת), כיוון שרינבוק יכול לאפשר לזיהום לחזור. ספר לרופא שלך אם יש לך פריחה כואבת על העור עם שלפוחיות, כיוון שאלו יכולים להיות סימנים לשלבקת חוגרת
- הייתה לך אי פעם דלקת כבד (הפטיטיס) מסוג B או C
- קיבלת לאחרונה או שאתה עתיד לקבל חיסון – זאת כיוון שחיסונים חיים אינם מומלצים בזמן השימוש ברינבוק
- יש לך סרטן – מכיוון שהרופא שלך יצטרך להחליט אם אתה עדיין יכול לקבל רינבוק
- אתה בסיכון גבוה לפתח סרטן עור, הרופא שלך יכול להמליץ על פעולות מונעות, כגון בדיקות עור שגרתיות בזמן נטילת רינבוק. דבר עם הרופא שלך אם אתה מפתח נגע חדש או שינוי במראה העור באזור מסוים.
- מטופלים מסוימים שטופלו ברינבוק פיתחו סרטני עור
- יש לך בעיות לב, לחץ דם גבוה או כולסטרול גבוה
- הכבד שלך אינו מתפקד כמו שצריך
- היו לך קרישי דם בוורידים הרגליים (פקקת ורידים עמוקים) או בריאות (תסחיף ריאתי). ספר לרופא שלך אם יש לך רגל נפוחה וכואבת, כאב בחזה או קוצר נשימה, כיוון שאלו יכולים להיות סימנים לקרישי דם בוורידים
- יש לך בעיות בכליות
- יש לך דלקת סעיף (דיברטיקוליטיס, סוג של דלקת במעי הגס) (ראה סעיף 4)

אם אתה חווה אחת מתופעות הלוואי הבאות, פנה מיד לרופא שלך:

- כאבי בטן עזים, במיוחד אלו המלווים בחום, בחילות והקאות

### בדיקות ומעקב

תצטרך לבצע בדיקות דם לפני התחלת הטיפול ברינבוק או בזמן נטילת התרופה. זאת כדי לבדוק אם קיימת ספירת תאי דם אדומים נמוכה (אנמיה), ספירת תאי דם לבנים נמוכה (נויטרופניה או לימפופניה), רמת שומנים גבוהה בדם (כולסטרול) או רמות גבוהות של אנזימי כבד. הבדיקות הן כדי לוודא שהטיפול ברינבוק אינו גורם לבעיות.

### קשישים

קיים שיעור גבוה יותר של זיהומים בקרב חולים בני 65 ומעלה. ספר לרופא שלך ברגע שאתה מבחין בסימנים או בתסמינים של זיהום.

### ילדים ומתבגרים

רינבוק לא מומלץ לשימוש בילדים מתחת לגיל 12 או במתבגרים השוקלים פחות מ-30 ק"ג הסובלים מדלקת עור אטופית, מכיוון שהוא לא נחקר בחולים אלו. רינבוק לא מומלץ לשימוש בילדים ובמתבגרים מתחת לגיל 18 עם דלקת מפרקים שגרונתית, דלקת מפרקים ספחתית או דלקת חוליות מקשחת. או קוליטיס כיבית, זאת מכיוון שהוא לא נחקר בקבוצת גיל זו.

### אינטראקציות/תגובות בין תרופתיות

**אם אתה לוקח, או אם לקחת לאחרונה, תרופות אחרות כולל תרופות ללא מרשם ותוספי תזונה, ספר על כך לרופא או לרוקח.** זאת מכיוון שתרופות מסוימות יכולות להפחית את יעילות הטיפול ברינבוק או עלולות להגביר את הסיכון לתופעות לוואי.

- תרופות לטיפול בזיהומים פטרייתיים (כגון איטראקונאזול, פוסאקונאזול או ווריקונאזול)
- תרופות לטיפול בזיהומים חיידקיים (כגון קלריתרומיצין)
- תרופות לטיפול בתסמונת קושינג (כגון קטוקונאזול)
- תרופות לטיפול בשחפת (כגון ריפאמפיצין)
- תרופות לטיפול בפרכוסים או התקפים (כגון פניטואין)
- תרופות המשפיעות על מערכת החיסון שלך (כגון אזאתיופרין, 6-מרקפטופורין, ציקלוספורין וטאקרולימוס)
- תרופות העלולות להעלות את הסיכון לדלקת סעיף (דיברטיקוליטיס) כמו תרופות נוגדות דלקת לא סטרואידיות (לרוב משמשות לטיפול בכאב ו/או מצבי דלקת של השרירים או המפרקים), ו/או אופיאטים (משמשים לטיפול בכאבים עזים), ו/או קורטיקוסטרואידים (לרוב משמשים לטיפול במצבי דלקת) (ראה סעיף 4)

אם אתה לוקח את אחת מהתרופות מהרשימה מעלה או שאינך בטוח, ספר לרופא או לרוקח שלך לפני נטילת רינבוק.

### הריון, הנקה ופוריות

#### הריון

אין להשתמש ברינבוק בזמן הריון.

#### הנקה

אם את מניקה או מתכננת להניק, דברי עם הרופא שלך לפני לקיחת תרופה זו. אין להשתמש ברינבוק בזמן הנקה כיוון שלא ידוע אם תרופה זו עוברת לחלב אם. את והרופא שלך תחליטו אם תניקי או תשתמשי ברינבוק. אל תעשי את שניהם.

#### מניעת הריון

אם את אישה בגיל הפוריות, עלייך להשתמש באמצעי מניעה יעילים כדי להימנע מהריון בזמן נטילת רינבוק ולמשך 4 שבועות לפחות לאחר מנת רינבוק האחרונה שלך. אם הרית בפרק זמן זה, עלייך לפנות מיד לרופא שלך.

אם הילדה שלך קיבלה לראשונה את המחזור החודשי ברינבוק, עליך לעדכן את הרופא.

### נהיגה ושימוש במכוונות

לרינבוק אין השפעה או השפעה מוגבלת על היכולת לנהוג ולהשתמש במכוונות.

### 3. כיצד תשתמש בתרופה ?

יש להשתמש בתכשיר תמיד בהתאם להוראות הרופא. עליך לבדוק עם הרופא או הרוקח אם אינך בטוח בנוגע למינון ואופן הטיפול בתכשיר.

המינון ואופן הטיפול יקבעו על ידי הרופא בלבד.

### **דלקת מפרקים שגרונית, דלקת מפרקים ספחתית ודלקת חוליות מקשחת**

המינון המקובל בדרך כלל הוא 15 מ"ג פעם אחת ביום.

### **דלקת עור אטופית**

**מבוגרים :**

המינון המקובל בדרך כלל הוא 15 מ"ג או 30 מ"ג פעם אחת ביום, כפי שנרשם על ידי הרופא שלך. הרופא עשוי להחליט אם להעלות או להוריד את המינון שלך בהתאם לפעולת התרופה בגוף.

**קשישים :**

אם אתה בן 65 ומעלה, המינון המקובל בדרך כלל הוא 15 מ"ג פעם אחת ביום.

**מתבגרים (גילאי 12-17) השוקלים לפחות 30 ק"ג :**

המינון המקובל בדרך כלל הוא 15 מ"ג פעם אחת ביום.

### **קוליטיס כיבית**

המינון המקובל בדרך כלל הוא 45 מ"ג פעם ביום למשך 8 שבועות. הרופא שלך עשוי להחליט להאריך את המינון ההתחלתי של 45 מ"ג למשך 8 שבועות נוספים (סה"כ 16 שבועות). לאחר מכן המינון המקובל בדרך כלל הוא 15 מ"ג או 30 מ"ג פעם ביום כטיפול ארוך טווח. הרופא עשוי להחליט להעלות או להוריד את המינון בהתאם לאופן שבו אתה מגיב לתרופה.

**קשישים :**

אם אתה בן 65 ומעלה, המינון המקובל בדרך כלל הוא 15 מ"ג פעם ביום כטיפול ארוך טווח.

הרופא עשוי להחליט להוריד את המינון אם יש לך בעיות בכליות, או שנרשמו לך תרופות מסוימות אחרות.

### **אין לעבור על המנה המומלצת**

#### **צורת הנטילה**

- יש לבלוע את הטבליה בשלמותה עם מים. אין לחצות, לכתוש, ללעוס או לשבור את הטבליה לפני הבליעה כיוון שזה יכול לשנות את כמות התרופה הנכנסת לגוף
- כדי לעזור לך לזכור לקחת רינבוק, קח אותו כל יום באותו הזמן
- ניתן לקחת את הטבליה עם או ללא מזון
- הימנע ממזון או משקה המכילים אשכוליות בזמן שאתה נוטל (או מטופל ב) רינבוק שכן אלו עלולים להגדיל את הסיכוי להופעת תופעות לוואי, על ידי הגדלת כמות התרופה בגוף

**אם נטלת בטעות מינון גבוה יותר** עליך לפנות לרופא שלך. ייתכן שתחווה מספר תופעות לוואי המפורטות בסעיף 4. אם נטלת מנת יתר או אם בטעות בלע ילד מן התרופה, פנה מיד לרופא או לחדר מיון של בית חולים והבא אריזת התרופה איתך.

## אם שכחת ליטול את התרופה

- אם פספסת מנה אחת, קח אותה מיד כשזוכרת
- אם שכחת את המנה שלך במשך יום שלם, דלג על המנה שהפסדת וקח רק מנה אחת כרגיל ביום למחרת
- אין לקחת מנה כפולה כדי לפצות על טבליה שנשכחה

יש להתמיד בטיפול כפי שהומלץ על ידי הרופא.

## אם אתה מפסיק את נטילת התרופה

אל תפסיק לקחת רינבוק אלא אם הרופא שלך הורה לך להפסיק לקחת אותו.

אין ליטול תרופות בחושך! בדוק התווית והמנה בכל פעם שהינך נוטל תרופה. הרכב משקפיים אם הינך זקוק להם. אם יש לך שאלות נוספות בנוגע לשימוש בתרופה זו, היוועץ ברופא או ברוקח.

## 4. תופעות לוואי

כמו בכל תרופה, השימוש ברינבוק עלול לגרום לתופעות לוואי בחלק מהמשתמשים. אל תיבהל למקרא רשימת תופעות הלוואי. ייתכן ולא תסבול מאף אחת מהן.

### תופעות לוואי חמורות

- דבר עם הרופא שלך או פנה מיד לקבלת עזרה רפואית אם אתה חווה סימנים כלשהם של זיהום, כגון:
- שלבכת חוגרת או פריחה כואבת על העור עם שלפוחיות (הרפס זוסטר) - תופעות לוואי שכיחות (common)
  - תופעות שמופיעות ב-10-1 משתמשים מתוך 100
  - זיהום בריאות (דלקת ריאות), אשר יכול לגרום לקוצר נשימה, חום ושיעול עם ליחה - תופעות לוואי שאינן שכיחות (uncommon) - תופעות שמופיעות ב-10-1 משתמשים מתוך 1,000

### תופעות לוואי אחרות

דבר עם הרופא שלך אם הבחנת באחת מתופעות הלוואי הבאות:

### תופעות לוואי שכיחות מאוד (very common) - תופעות שמופיעות ביותר ממשתמש אחד מעשרה

- זיהומים בגרון ובאף
- אקנה

### תופעות לוואי שכיחות (common) - תופעות שמופיעות ב-10-1 משתמשים מתוך 100

- שיעול
- חום
- פצעי קור (הרפס סימפלקס)
- בחילה
- עלייה ברמת אנזים אשר נקרא קראטין קינאז, הנראית בבדיקות דם
- ספירת תאי דם לבנים נמוכה, הנראית בבדיקות דם
- עלייה ברמות כולסטרול (סוג של שומן בדם) כפי שנראה בבדיקות
- עלייה ברמות אנזימי הכבד, הנראית בבדיקות דם (סימן לבעיות כבד)
- עלייה במשקל
- דלקת (נפיחות) של זקיקי השיער
- שפעת
- אנמיה
- כאבים בבטן

- עייפות (תחושת עייפות בלתי רגילה וחולשה)
- כאב ראש
- סרפדת (אורטיקריה)
- זיהום בדרכי השתן
- **פריחה**

**תופעות לוואי שאינן שכיחות (uncommon) - תופעות שמופיעות ב-10-1 משתמשים מתוך 1,000**

- פטרת הפה (כתמים לבנים בפה)
- עלייה ברמות טריגליצרידים (סוג של שומן) בדם כפי שנראה בבדיקות
- דלקת הסעיף (דיברטיקוליטיס, התפתחות דלקת כואבת בכיסים קטנים שנוצרים בדופן המעי שלך)

אם הופיעה תופעת לוואי, אם אחת מתופעות הלוואי מחמירה או כאשר אתה סובל מתופעת לוואי שלא צוינה בעלון, עליך להתייעץ עם הרופא.

**דיווח תופעות לוואי**

ניתן לדווח על תופעות לוואי למשרד הבריאות באמצעות לחיצה על הקישור "דיווח על תופעות לוואי עקב טיפול תרופתי" שנמצא בדף הבית של אתר משרד הבריאות ([www.health.gov.il](http://www.health.gov.il)) המפנה לטופס המקוון לדיווח על תופעות לוואי, או על ידי כניסה לקישור:

<https://sideeffects.health.gov.il/>

**5. איך לאחסן את התרופה?**

- מנע הרעלה! תרופה זו וכל תרופה אחרת יש לשמור במקום סגור מחוץ להישג ידם וטווח ראייתם של ילדים ו/או תינוקות ועל ידי כך תמנע הרעלה. אל תגרום להקאה ללא הוראה מפורשת מהרופא
- אין להשתמש בתרופה אחרי תאריך התפוגה (exp. date) המופיע על גבי האריזה החיצונית והבליסטר. תאריך התפוגה מתייחס ליום האחרון של אותו חודש
- יש לאחסן בטמפרטורה עד 30°C
- יש לאחסן בבליסטר המקורי כדי להגן מלחות
- אין להשליך תרופות כלשהן דרך הביוב או האשפה הביתית. שאל את הרוקח שלך כיצד להשליך תרופות אשר אינך משתמש בהן יותר. אמצעים אלו יעזרו לשמור על הסביבה

**6. מידע נוסף**

**מה מכיל רינבוק**

נוסף על המרכיב הפעיל התרופה מכילה גם:

רינבוק 15 מ"ג טבליות לשחרור ממושך

- ליבת הטבליה:

microcrystalline cellulose, mannitol, hypromellose, tartaric acid (powdered), magnesium stearate, silica colloidal anhydrous/ colloidal silicon dioxide

- ציפוי הפילם:

polyvinyl alcohol, macrogol/polyethylene glycol, talc, titanium dioxide (E171), black iron oxide (E172)/ferrosoferric oxide, iron oxide red (E172)

רינבוק 30 מ"ג טבליות לשחרור ממושך

- ליבת הטבליה:

microcrystalline cellulose, mannitol, hypromellose, tartaric acid (powdered), magnesium stearate, silica colloidal anhydrous/ colloidal silicon dioxide

● ציפוי הפילם :

polyvinyl alcohol, macrogol/ polyethylene glycol, talc, titanium dioxide (E171), iron oxide red (E172).

### **כיצד נראית התרופה ומה תוכן האריזה**

רינבוק 15 מ"ג טבליות לשחרור ממושך

רינבוק 15 מ"ג טבליות לשחרור ממושך הינן סגולות, מוארכות, קמורות משני הצדדים, מודפסות על צד אחד עם 'a15'.

הטבליות מסופקות בבליסטרים באריזות המכילות 28 או 98 טבליות לשחרור ממושך ובמארז של 84 המכיל 3 קופסאות, כל אחת מכילה 28 טבליות לשחרור ממושך. כל בליסטר המציג את ימי השבוע מכיל 7 טבליות.

רינבוק 30 מ"ג טבליות לשחרור ממושך

רינבוק 30 מ"ג טבליות לשחרור ממושך הינן אדומות, מוארכות, קמורות משני הצדדים, מודפסות על צד אחד עם 'a30'.

הטבליות מסופקות בבליסטרים באריזות המכילות 28 או 98 טבליות לשחרור ממושך. כל בליסטר המציג את ימי השבוע מכיל 7 טבליות.

ייתכן ולא כל גודלי האריזות משווקים.

● **בעל הרישום וכתובתו:** [AbbVie Biopharmaceuticals Ltd.](http://AbbVie-Biopharmaceuticals-Ltd), רח' החרש 4, הוד השרון, ישראל

● **שם היצרן וכתובתו:** [AbbVie Deutschland](http://AbbVie-Deutschland), [GmbH & Co. KG](http://GmbH-&Co.-KG), Knollstrasse 67061, לודויגשפן, גרמניה

● נערך ב**יולי-באוגוסט** 2022 בהתאם להנחיות משרד הבריאות

● **מספר רישום התרופה בפנקס התרופות הממלכתי במשרד הבריאות:**  
רינבוק 15 מ"ג טבליות לשחרור ממושך 164-29-36194  
רינבוק 30 מ"ג טבליות לשחרור ממושך 168-69-37036



למידע נוסף ולמוקד תוכנית התמיכה

התקשרו \*6718

לשם הפשטות ולהקלת הקריאה, עלון זה נוסח בלשון זכר. על אף זאת, התרופה מיועדת לבני שני המינים.